

METAL-BINDING COMPOUNDS AND USES THEREFOR

This application claims benefit of provisional applications 60/157,404, filed October 1, 1999, and 60/211,078, filed June 13, 2000.

5

FIELD OF THE INVENTION

10

The invention relates to a method of reducing the molecular, cellular and tissue damage done by reactive oxygen species (ROS). The invention also relates to certain compounds, especially certain peptides and peptide derivatives, that bind metal ions, particularly Cu(II). The binding of metal ions by the compounds of the invention inhibits the formation and/or accumulation of ROS and/or targets the damage done by ROS to the compounds themselves (*i.e.*, the compounds of the invention may act as sacrificial antioxidants).

15

BACKGROUND

20

25

Reactive oxygen species (ROS) include free radicals (*e.g.*, superoxide anion and hydroxyl, peroxy, and alkoxy radicals) and non-radical species (*e.g.*, singlet oxygen and hydrogen peroxide). ROS are capable of causing extensive cellular and tissue damage, and they have been reported to play a major role in a variety of diseases and conditions. Indeed, ROS have been implicated in over 100 diseases and pathogenic conditions, and it has been speculated that ROS may constitute a common pathogenic mechanism involved in all human diseases. Stohs, *J. Basic Clin. Physiol. Pharmacol.*, **6**, 205-228 (1995). For reviews describing ROS, their formation, the mechanisms by which they cause cellular and tissue damage, and their involvement in numerous diseases and disorders, see, *e.g.*, Manso, *Rev. Port. Cardiol.*, **11**, 997-999 (1992); Florence, *Aust. NZ J. Ophthalmol.*, **23**, 3-7 (1992); Stohs, *J. Basic Clin. Physiol. Pharmacol.*, **6**, 205-228 (1995); Knight, *Ann. Clin. Lab. Sci.*, **25**, 111-121 (1995); Kerr et al., *Heart & Lung*, **25**, 200-209 (1996); Roth, *Acta Chir. Hung.*, **36**, 302-305 (1997).

Ischemia/reperfusion is the leading cause of illness and disability in the world. Cardiovascular ischemia, in which the body's capacity to provide oxygen to the heart is diminished, is the leading cause of illness and death in the United States. Cerebral ischemia is a precursor to cerebrovascular accident (stroke), which is the third leading cause of death in the United States. Ischemia also occurs in other organs (*e.g.*, kidney, liver, lung, and the intestinal tract), in harvested organs (*e.g.*, organs harvested for transplantation or for research (*e.g.*, perfused organ models)), and as a result of surgery where blood flow is interrupted (*e.g.*, open heart surgery and coronary bypass surgery). Ischemia need not be limited to one organ; it can also be more generalized (*e.g.*, in hemorrhagic shock).

Cellular and tissue damage occur during ischemia as result of oxygen deficiency. However, the damage that occurs during ischemia is generally light compared to the severe damage that occurs upon reperfusion of ischemic tissues and organs. See, *e.g.*, Manso, *Rev. Port. Cardiol.*, **11**, 997-999 (1992); Stohs, *J. Basic Clin. Physiol. Pharmacol.*, **6**, 205-228 (1995); Knight, *Ann. Clin. Lab. Sci.*, **25**, 111-121 (1995); Kerr et al., *Heart & Lung*, **25**, 200-209 (1996); Roth, *Acta Chir. Hung.*, **36**, 302-305 (1997). ROS have been reported to be responsible for the severe damage caused by reperfusion of ischemic tissues and organs. See, *e.g.*, Manso, *Rev. Port. Cardiol.*, **11**, 997-999 (1992); Stohs, *J. Basic Clin. Physiol. Pharmacol.*, **6**, 205-228 (1995); Knight, *Ann. Clin. Lab. Sci.*, **25**, 111-121 (1995); Kerr et al., *Heart & Lung*, **25**, 200-209 (1996); Roth, *Acta Chir. Hung.*, **36**, 302-305 (1997).

Metal ions, primarily transition metal ions, can cause the production and accumulation of ROS. In particular, copper and iron ions released from storage sites are one of the main causes of the production of ROS following injury, including ischemia/reperfusion injury and injury due to heat, cold, trauma, excess exercise, toxins, radiation, and infection. Roth, *Acta Chir. Hung.*, **36**, 302-305 (1997). Copper and iron ions, as well as other transition metal ions (*e.g.*, vanadium, and chromium ions), have been reported to catalyze the production of ROS. See, *e.g.*, Stohs, *J. Basic Clin. Physiol. Pharmacol.*, **6**, 205-228 (1995); Halliwell et al., *Free Radicals In Biology And Medicine*, pages 1-19 (Oxford University 1989); Marx et al., *Biochem. J.*, **236**, 397-400 (1985); Quinlan et al., *J. Pharmaceutical Sci.*, **81**, 611-614

(1992). Other transition metal ions (*e.g.*, cadmium, mercury, and nickel ions) and other metal ions (*e.g.*, arsenic and lead ions) have been reported to deplete some of the molecules of the natural antioxidant defense system, thereby causing an increased accumulation of ROS. *See, e.g.*, Stohs, *J. Basic Clin. Physiol. Pharmacol.*, **6**, 205-228 (1995). Although it has been reported that free copper ions bind nonspecifically to the amino groups of essentially any protein (Gutteridge et al., *Biochim. Biophys. Acta*, **759**, 38-41 (1983)), copper ions bound to proteins can still cause the production of ROS which damage at least the protein to which the copper ions are bound. *See, e.g.*, Gutteridge et al., *Biochim. Biophys. Acta*, **759**, 38-41 (1983); Marx et al., *Biochem. J.*, **236**, 397-400 (1985); Quinlan et al., *J. Pharmaceutical Sci.*, **81**, 611-614 (1992).

Albumin has been characterized as an extracellular antioxidant. *See, e.g.*, Halliwell and Gutteridge, *Arch. Biochem. Biophys.*, **280**, 1-8 (1990); Das et al., *Methods Enzymol.*, **233**, 601-610 (1994); Stohs, *J. Basic Clin. Physiol. Pharmacol.*, **6**, 205-228 (1995); Dunphy et al., *Am. J. Physiol.*, **276**, H1591-H1598 (1999)). The antioxidant character of albumin has been attributed to several of albumin's many physiological functions, including albumin's ability to bind metals (particularly copper ions), to bind fatty acids, to bind and transport steroids, to bind and transport bilirubin, to scavenge HOCl, and others. *See, e.g.*, Halliwell and Gutteridge, *Arch. Biochem. Biophys.*, **280**, 1-8 (1990); Halliwell and Gutteridge, *Arch. Biochem. Biophys.*, **246**, 501-514 (1986); Stohs, *J. Basic Clin. Physiol. Pharmacol.*, **6**, 205-228 (1995); Dunphy et al., *Am. J. Physiol.*, **276**, H1591-H1598 (1999)). Albumin contains several metal binding sites, including one at the N-terminus. The N-terminal metal-binding sites of several albumins, including human, rat and bovine serum albumins, exhibit high-affinity for Cu(II) and Ni(II), and the amino acids involved in the high-affinity binding of these metal ions have been identified. *See* Laussac et al., *Biochem.*, **23**, 2832-2838 (1984); Predki et al., *Biochem. J.*, **287**, 211-215 (1992); Masuoka et al., *J. Biol. Chem.*, **268**, 21533-21537 (1993). It has been reported that copper bound to albumin at metal binding sites other than the high-affinity N-terminal site produce free radicals which causes extensive damage to

albumin at sites dictated by the location of the “loose” metal binding sites, resulting in the characterization of albumin as a “sacrificial antioxidant.” See Marx et al., *Biochem. J.*, **236**, 397-400 (1985); Halliwell et al., *Free Radicals In Biology And Medicine*, pages 1-19 (Oxford University 1989); Halliwell and Gutteridge, *Arch. Biochem. Biophys.*, **280**, 1-8 (1990);
 5 Quinlan et al., *J. Pharmaceutical Sci.*, **81**, 611-614 (1992).

Despite the foregoing, attempts to use albumin as a treatment for cerebral ischemia have shown mixed results. It has been reported that albumin is, and is not, neuroprotective in animal models of cerebral ischemia. Compare Huh et al., *Brain Res.*, **804**, 105-113 (1998) and Remmers et al., *Brain Res.*, **827**, 237-242 (1999), with Little et al., *Neurosurgery*, **9**,
 10 552-558 (1981) and Beaulieu et al., *J. Cereb. Blood Flow. Metab.*, **18**, 1022-1031 (1998).

Mixed results have also been obtained using albumin in cardioplegia solutions for the preservation of excised hearts. As reported in Dunphy et al., *Am. J. Physiol.*, **276**, H1591-H1598 (1999), the addition of albumin to a standard cardioplegia solution for the preservation of excised hearts did not improve the functioning of hearts perfused with the solution for twenty-four hours. Hearts did demonstrate improved functioning when perfused with a cardioplegia solution containing albumin and several enhancers (insulin, ATP, corticosterone, and pyruvic acid). This was a synergistic effect, since the enhancers alone, as well as the albumin alone, did not significantly improve heart function. An earlier report of improved heart function using cardioplegia solutions containing albumin was also attributed to synergism between enhancers and albumin. See the final paragraph of Dunphy et al., *Am. J. Physiol.*, **276**, H1591-H1598 (1999) and Hisatomi et al., *Transplantation*, **52**, 754-755 (1991), cited therein. In another study, hearts perfused with a cardioplegia solution containing albumin increased reperfusion injury in a dose-related manner, as compared to a solution not containing albumin. Suzer et al., *Pharmacol. Res.*, **37**, 97-101 (1998). Based
 25 on their study and the studies of others, Suzer et al. concluded that albumin had not been shown to be effective for cardioprotection. They further noted that the use of albumin in cardioplegia solutions could be unsafe due to possible allergic reactions and the risks associated with the use of blood products.

Finally, although albumin has been characterized as an antioxidant, it has also been reported to enhance superoxide anion production by microglia (Si et al., *GLIA*, **21**, 413-418 (1997)). This result led the authors to speculate that albumin leaking through the disrupted blood brain barrier in certain disorders potentiates the production of superoxide anion by microglia, and that this increased production of superoxide anion is responsible for the pathogenesis of neuronal damage in cerebral ischemia/reperfusion and some neurodegenerative diseases.

As noted above, the N-terminal metal-binding sites of several albumins exhibit high-affinity for Cu(II) and Ni(II). These sites have been studied extensively, and a general amino terminal Cu(II)- and Ni(II)-binding (ATCUN) motif has been identified. See, *e.g.*, Harford and Sarkar, *Acc. Chem. Res.*, **30**, 123-130 (1997). The ATCUN motif can be defined as being present in a protein or peptide which has a free -NH₂ at the N-terminus, a histidine residue in the third position, and two intervening peptide nitrogens. See, *e.g.*, Harford and Sarkar, *Acc. Chem. Res.*, **30**, 123-130 (1997). Thus, the ATCUN motif is provided by the peptide sequence Xaa Xaa His, where Xaa is any amino acid except proline. See, *e.g.*, Harford and Sarkar, *Acc. Chem. Res.*, **30**, 123-130 (1997). The Cu(II) and Ni(II) are bound by four nitrogens provided by the three amino acids of the ATCUN motif (the nitrogen of the free -NH₂, the two peptide nitrogens, and an imidazole nitrogen of histidine) in a slightly distorted square planar configuration. See, *e.g.*, Harford and Sarkar, *Acc. Chem. Res.*, **30**, 123-130 (1997). Side-chain groups of the three amino acids of which the ATCUN motif consists can be involved in the binding of the Cu(II) and Ni(II), and amino acids near these three N-terminal amino acids may also have an influence on the binding of these metal ions. See, *e.g.*, Harford and Sarkar, *Acc. Chem. Res.*, **30**, 123-130 (1997); Bal et al., *Chem. Res. Toxicol.*, **10**, 906-914 (1997). For instance, the sequence of the N-terminal metal-binding site of human serum albumin is Asp Ala His Lys [SEQ ID NO:1], and the free side-chain carboxyl of the N-terminal Asp and the Lys residue have been reported to be involved in the binding of Cu(II) and Ni(II), in addition to the four nitrogens provided by Asp Ala His. See Harford

and Sarkar, *Acc. Chem. Res.*, **30**, 123-130 (1997); Laussac et al., *Biochem.*, **23**, 2832-2838 (1984); and Sadler et al., *Eur. J. Biochem.*, **220**, 193-200 (1994).

The ATCUN motif has been found in other naturally-occurring proteins besides albumins, and non-naturally-occurring peptides and proteins comprising the ATCUN motif have been synthesized. See, e.g., Harford and Sarkar, *Acc. Chem. Res.*, **30**, 123-130 (1997); Bal et al., *Chem. Res. Toxicol.*, **10**, 906-914 (1997); Mlynarz, et al., *Speciation 98: Abstracts*, <http://www.jate.u-szeged.hu/~spec98/abstr/mlynar.html>. Cu(II) and Ni(II) complexes of ATCUN-containing peptides and proteins have been reported to exhibit superoxide dismutase (SOD) activity. See Cotelle et al., *J. Inorg. Biochem.*, **46**, 7-15 (1992); Ueda et al., *J. Inorg. Biochem.*, **55**, 123-130 (1994). Despite their reported SOD activity, these complexes still produce free radicals which damage DNA, proteins and other biomolecules. See Harford and Sarkar, *Acc. Chem. Res.*, **30**, 123-130 (1997); Bal et al., *Chem. Res. Toxicol.*, **10**, 915-21 (1997); Ueda et al., *Free Radical Biol. Med.*, **18**, 929-933 (1995); Ueda et al., *J. Inorg. Biochem.*, **55**, 123-130 (1994); Cotelle et al., *J. Inorg. Biochem.*, **46**, 7-15 (1992). As a consequence, it has been hypothesized that at least some of the adverse effects of copper and nickel *in vivo* are attributable to the binding of Cu(II) and Ni(II) to ATCUN-containing proteins which causes the production of damaging free radicals. See Harford and Sarkar, *Acc. Chem. Res.*, **30**, 123-130 (1997); Bal et al., *Chem. Res. Toxicol.*, **10**, 915-921 (1997); Cotelle et al., *J. Inorg. Biochem.*, **46**, 7-15 (1992). Cf. Koch et al., *Chem. & Biol.*, **4**, 549-60 (1997). The damaging effects produced by a Cu(II) complex of an ATCUN-containing peptide have been exploited to kill cancer cells *in vitro* and to produce anti-tumor effects *in vivo*. See Harford and Sarkar, *Acc. Chem. Res.*, **30**, 123-130 (1997).

SUMMARY OF THE INVENTION

The invention provides a method of reducing the damage done by reactive oxygen species (ROS) in an animal. The method comprises administering to the animal an effective amount of a metal-binding peptide having the formula $P_1 - P_2$ or a physiologically-acceptable salt thereof.

The invention further provides a method of reducing the damage done by ROS in a tissue or an organ that has been removed from an animal. This method comprises contacting the tissue or organ with a solution containing an effective amount of the peptide $P_1 - P_2$ or a physiologically-acceptable salt thereof.

The invention also provides a method of reducing the concentration of a metal in an animal in need thereof. The method comprises administering to the animal an effective amount of a metal-binding peptide having the formula $P_1 - P_2$ or a physiologically-acceptable salt thereof.

The invention also provides a pharmaceutical composition comprising a pharmaceutically-acceptable carrier and the peptide $P_1 - P_2$ or a physiologically-acceptable salt thereof.

In addition, the invention provides a kit for reducing the damage done by ROS in a tissue or organ that has been removed from an animal. The kit comprises a container holding the peptide $P_1 - P_2$.

In the formula $P_1 - P_2$:

P_1 is $Xaa_1 Xaa_2 His$ or $Xaa_1 Xaa_2 His Xaa_3$; and

P_2 is $(Xaa_4)_n$.

Xaa_1 is glycine (Gly), alanine (Ala), valine (Val), leucine (Leu), isoleucine (Ile), serine (Ser), threonine (Thr), aspartic acid (Asp), asparagine (Asn), glutamic acid (Glu), glutamine (Gln), lysine (Lys), hydroxylysine (Hylys), histidine (His), arginine (Arg), ornithine (Orn), phenylalanine (Phe), tyrosine (Tyr), tryptophan (Trp), cysteine (Cys), methionine (Met) or α -hydroxymethylserine (HMS). Xaa_1 is preferably Asp, Glu, Arg, or HMS. More preferably, Xaa_1 is Asp or Glu. Most preferably Xaa_1 is Asp.

Xaa₂ is Gly, Ala, β -Ala, Val, Leu, Ile, Ser, Thr, Asp, Asn, Glu, Gln, Lys, Hylys, His, Arg, Orn, Phe, Tyr, Trp, Cys, Met or HMS. Xaa₂ is preferably Gly, Ala, Val, Leu, Ile, Thr, Ser, Asn, Met, His or HMS. More preferably Xaa₂ is Ala, Val, Thr, Ser, or HMS. Even more preferably Xaa₂ is Ala, Thr, or HMS. Most preferably Xaa₂ is Ala.

5 Xaa₃ is Gly, Ala, Val, Lys, Arg, Orn, Asp, Glu, Asn, Gln, or Trp, preferably Lys.

Xaa₄ is any amino acid.

Finally, n is 0-100, preferably 0-10, more preferably 0-5, and most preferably 0.

In a preferred embodiment, at least one of the amino acids of P₁, other than β -Ala when it is present, is a D-amino acid. Preferably, the D-amino acid is Xaa₁, His, or both.
 10 Most preferably all of the amino acids of P₁, other than β -Ala, are D-amino acids. In another preferred embodiment, at least one of the amino acids of P₁ other than β -Ala is a D-amino acid, and at least 50% of the amino acids of P₂ are also D-amino acids. Most preferably all of the amino acids of P₂ are D-amino acids.

In another preferred embodiment, at least one amino acid of P₁ and/or P₂ is substituted with (a) a substituent that increases the lipophilicity of the peptide without altering the ability of P₁ to bind metal ions, (b) a substituent that protects the peptide from proteolytic enzymes without altering the ability of P₁ to bind metal ions, or (c) a substituent which is a non-peptide, metal-binding functional group that improves the ability of the peptide to bind metal ions.
 15

20 The invention provides another method of reducing the damage done by ROS in an animal. The method comprises administering to the animal an effective amount of a metal-binding peptide (MBP) having attached thereto a non-peptide, metal-binding functional group. The metal-binding peptide MBP may be any metal-binding peptide, not just P₁ - P₂.

25 The invention further provides another method of reducing the damage done by ROS in a tissue or an organ that has been removed from an animal. This method comprises contacting the tissue or organ with a solution containing an effective amount of a metal-binding peptide MBP having attached thereto a non-peptide, metal-binding functional group.

The invention provides another method of reducing the concentration of a metal in an animal in need thereof. The method comprises administering to the animal an effective amount of a metal-binding peptide MBP having attached thereto a non-peptide, metal-binding functional group.

5 The invention also provides a pharmaceutical composition comprising a pharmaceutically-acceptable carrier and a metal-binding peptide MBP having attached thereto a non-peptide, metal-binding functional group.

The invention also provides a kit for reducing the damage done by ROS in a tissue or organ that has been removed from an animal. The kit comprises a container holding a
10 metal-binding peptide MBP having attached thereto a non-peptide, metal-binding functional group.

The invention provides yet another method of reducing the damage done by reactive oxygen species (ROS) in an animal. The method comprises administering to the animal an effective amount of a metal-binding peptide dimer of the formula $P_3 - L - P_3$, wherein each P_3 may be the same or different and is a peptide which is capable of binding a metal ion, and L is a chemical group which connects the two P_3 peptides through their C-terminal amino acids. In a preferred embodiment, one or both of the two P_3 peptides is P_1 .
15

The invention further provides a method of reducing the damage done by ROS in a tissue or an organ that has been removed from an animal. This method comprises contacting the tissue or organ with a solution containing an effective amount of the metal-binding peptide dimer of the formula $P_3 - L - P_3$.
20

The invention also provides a method of reducing the concentration of a metal in an animal in need thereof. The method comprises administering to the animal an effective amount of the metal-binding peptide dimer of the formula $P_3 - L - P_3$.
25

The invention also provides a pharmaceutical composition comprising a pharmaceutically-acceptable carrier and the metal-binding peptide dimer of the formula $P_3 - L - P_3$.

In addition, the invention provides a kit for reducing the damage done by ROS in a tissue or organ that has been removed from an animal. The kit comprises a container holding the metal-binding peptide dimer of the formula $P_3 - L - P_3$.

In addition, the invention provides a peptide having the formula $P_1 - P_2$, or a physiologically-acceptable salt thereof, wherein at least one amino acid of P_1 , other than β -Ala, is a D-amino acid.

Further provided by the invention is a peptide having the formula $P_1 - P_2$, or a physiologically-acceptable salt thereof, wherein at least one amino acid of P_1 and/or P_2 is substituted with (a) a substituent that increases the lipophilicity of the peptide without altering the ability of P_1 to bind metal ions, (b) a substituent that protects the peptide from proteolytic enzymes without altering the ability of P_1 to bind metal ions, or (c) a substituent which is a non-peptide, metal-binding functional group that improves the ability of the peptide to bind metal ions.

In addition, the invention provides a peptide having the formula $P_1 - P_2$, wherein P_1 is defined above, and P_2 is a peptide sequence which comprises the sequence of a metal-binding site. In particular, P_2 may have one of the following sequences: $(Xaa_4)_m Xaa_3 His Xaa_2 Xaa_5$, $(Xaa_4)_m His Xaa_2 Xaa_5$, $(Xaa_4)_m Xaa_5 Xaa_2 His Xaa_3$, or $(Xaa_4)_m Xaa_5 Xaa_2 His$, where Xaa_5 is an amino acid having a free side-chain $-NH_2$ and m is 0-5.

The invention also provides a metal-binding peptide MBP having attached thereto a non-peptide, metal-binding functional group.

Finally, the invention provides the metal-binding peptide dimer of the formula $P_3 - L - P_3$.

BRIEF DESCRIPTION OF THE DRAWINGS

Figures 1A-D: Formulas of tetrapeptide Asp Ala His Lys [SEQ ID NO:1] showing points of possible substitution.

Figures 2A-B: Schematic diagrams of the synthesis of derivatives of the tetrapeptide Asp Ala His Lys [SEQ ID NO:1] coming within the formula of Figure 1C (Figure 2A) and Figure 1B (Figure 2B).

Figure 3A-B: Formulas of cyclohexane diamine derivatives.

Figures 3C-D: Schematic diagrams of syntheses of cyclohexane diamine derivatives of the tetrapeptide Asp Ala His Lys [SEQ ID NO:1].

Figure 4: Formula of a tetraacetic acid derivative of the tetrapeptide Asp Ala His Lys [SEQ ID NO:1].

Figure 5: Formula of a bispyridylethylamine derivative of the tetrapeptide Asp Ala His Lys [SEQ ID NO:1].

Figures 6A-B: Formulas of mesoporphyrin IX with (Figure 6B) and without (Figure 6A) a bound metal ion M.

Figure 6C: Formula of mesoporphyrin IX derivative of the tetrapeptide Asp Ala His Lys [SEQ ID NO:1].

Figure 7: Formulas of monosaccharides.

Figure 8: Diagram of a parabiologic blood perfusion system in which an isolated heart is perfused in the Langendorff mode with blood at 37°C from a support animal of the same species.

Figure 9: Diagram of the treatments of isolated perfused hearts with drug and saline in the parabiologic blood perfusion system illustrated in Figure 8.

Figure 10: Graph of contracture versus duration of ischemia showing the effect of a drug (D-Asp D-Ala D-His D-Lys) on contracture during ischemia in the blood-perfused rat heart model illustrated in Figures 8 and 9. In Figure 10, -□- is saline control, and -○- is drug.

Figure 11: Graph of left ventricle diastolic pressure (LVDP; expressed as a percentage of the 20-minute pre-intervention baseline value) versus duration of reperfusion showing the effect of the drug D-Asp D-Ala D-His D-Lys on post-ischemic recovery of LVDP in the blood-perfused rat heart model illustrated in Figures 8 and 9. * indicates $p \leq 0.05$. In Figure 11, -□- is saline control, and -○- is drug.

Figure 12: Graph of left ventricle end diastolic pressure (LVEDP) versus duration of reperfusion showing the effect of the drug D-Asp D-Ala D-His D-Lys on post-ischemic recovery of LVEDP in the blood-perfused rat heart model illustrated in Figures 8 and 9. * indicates $p \leq 0.05$. In Figure 12, -□- is saline control, and -○- is drug.

Figure 13: Graph of heart rate (expressed as a percentage of the 20-minute pre-intervention baseline value) versus duration of reperfusion showing the effect of the drug D-Asp D-Ala D-His D-Lys on post-ischemic recovery of heart rate in the blood-perfused rat heart model illustrated in Figures 8 and 9. * indicates $p \leq 0.05$. In Figure 13, -□- is saline control, and -○- is drug.

Figure 14: Graph of perfusion pressure (expressed as a percentage of the 20-minute pre-intervention baseline value) versus duration of reperfusion showing the effect of the drug D-Asp D-Ala D-His D-Lys on post-ischemic recovery of perfusion pressure in the blood-perfused rat heart model illustrated in Figures 8 and 9. In Figure 14, -□- is saline control, and -○- is drug.

Figure 15A-B: Graphs of absorbance at 532 nm (A532) versus incubation time in an assay for the production of hydroxyl radicals. In Figure 15A, ■ = ascorbate only, ◆ = copper and ascorbate, ▲ = tetrapeptide (L-Asp L-Ala L-His L-Lys [SEQ ID NO:1]), copper and ascorbate (tetrapeptide/copper ratio of 1:1), X = tetrapeptide, copper and ascorbate (tetrapeptide/copper ratio of 2:1). In Figure 15B, ◆ = copper and ascorbate and ■ = tetrapeptide, copper and ascorbate (tetrapeptide/copper ratio of 2:1).

Figure 16: Graph of % inhibition versus concentration tetrapeptide (L-Asp L-Ala L-His L-Lys [SEQ ID NO:1])-copper complex at a tetrapeptide/copper ratio of 1:1 in the xanthine oxidase assay for superoxide dismutase activity.

Figure 17: Graph of absorbance at 560 nm (A560) versus time in an assay for superoxide radical production. In Figure 17, ■ = ascorbate only, ◆ = copper and ascorbate, Δ = tetrapeptide (L-Asp L-Ala L-His L-Lys [SEQ ID NO:1]), copper and ascorbate (tetrapeptide/copper ratio of 1:1), X = tetrapeptide, copper and ascorbate (tetrapeptide/copper ratio of 2:1).

Figure 18: Gel after electrophoresis of DNA treated in various ways. Lane 1 - 17 $\mu\text{g/ml}$ plasmid DNA (untreated control); Lane 2 - 17 $\mu\text{g/ml}$ plasmid DNA and 50 μM CuCl_2 ; Lane 3 - 17 $\mu\text{g/ml}$ plasmid DNA and 2.5 mM ascorbate; Lane 4 - 17 $\mu\text{g/ml}$ plasmid DNA, 2.5 mM ascorbate, 50 μM CuCl_2 , and 200 μM tetrapeptide (L-Asp L-Ala L-His L-Lys [SEQ ID NO:1]) (4:1 ratio tetrapeptide/copper); Lane 5 - 17 $\mu\text{g/ml}$ plasmid DNA, 2.5 mM ascorbate, 50 μM CuCl_2 , and 100 μM tetrapeptide (2:1 ratio tetrapeptide/copper); Lane 6 - 17 $\mu\text{g/ml}$ plasmid DNA, 2.5 mM ascorbate, 50 μM CuCl_2 , and 50 μM tetrapeptide (1:1 ratio tetrapeptide/copper); Lane 7 - 17 $\mu\text{g/ml}$ plasmid DNA, 2.5 mM ascorbate, 50 μM CuCl_2 , and 25 μM tetrapeptide (1:2 ratio tetrapeptide/copper); Lane 8 - 17 $\mu\text{g/ml}$ plasmid DNA, 2.5 mM ascorbate, 50 μM CuCl_2 , and 12.5 μM tetrapeptide (1:4 ratio tetrapeptide/copper); Lane 9 - 17 $\mu\text{g/ml}$ plasmid DNA, 2.5 mM ascorbate, and 50 μM CuCl_2 (positive control); and Lane 10 - DNA ladder.

Figure 19A: Formulas of peptide dimers according to the invention.

Figures 19B-C: Diagrams illustrating the synthesis of peptide dimers according to the invention.

DETAILED DESCRIPTION OF THE PRESENTLY-PREFERRED EMBODIMENTS

The invention provides a peptide of the formula $P_1 - P_2$. P_1 is $\text{Xaa}_1 \text{Xaa}_2 \text{His}$ or is $\text{Xaa}_1 \text{Xaa}_2 \text{His Xaa}_3$, wherein Xaa_1 , Xaa_2 , and Xaa_3 are defined above. P_1 is a metal-binding peptide sequence that binds transition metal ions of Groups 1b-7b or 8 of the Periodic Table of elements (including V, Co, Cr, Mo, Mn, Ba, Zn, Hg, Cd, Au, Ag, Co, Fe, Ni, and Cu) and other metal ions (including As, Sb and Pb). The binding of metal ions by P_1 inhibits (*i.e.*, reduces or prevents) the production of ROS and/or the accumulation of ROS by these metal ions and/or targets the damage done by ROS that may still be produced by the bound metal ions to the peptide itself. As a result, the damage that can be caused by ROS in the absence of the binding of the metal ions to P_1 is reduced. In particular, P_1 binds Cu(II), Ni(II), Co(II),

and Mn(II) with high affinity. It should, therefore, be particularly effective in reducing the damage caused by the production and accumulation of ROS by copper and nickel.

In P_1 , Xaa_1 is most preferably Asp, Xaa_2 is most preferably Ala, and Xaa_3 is most preferably Lys. Thus, the preferred sequences of P_1 are Asp Ala His and Asp Ala His Lys [SEQ ID NO:1]. Most preferably the sequence of P_1 is Asp Ala His Lys [SEQ ID NO:1]. Asp Ala His is the minimum sequence of the N-terminal metal-binding site of human serum albumin necessary for the high-affinity binding of Cu(II) and Ni(II), and Lys has been reported to contribute to the binding of these metal ions to this site. Other sequences for P_1 may be preferred for use in animals other than humans.

P_2 is $(Xaa_4)_n$, wherein Xaa_4 is any amino acid and n is 0-100. When n is large ($n >$ about 20), the peptides will reduce the damage done by ROS extracellularly. Smaller peptides are better able to enter cells, and smaller peptides can, therefore, be used to reduce the damage done by ROS both intracellularly and extracellularly. Smaller peptides are also less subject to proteolysis. Therefore, in P_2 , preferably n is 0-10, more preferably n is 0-5, and most preferably n is 0. Although P_2 may have any sequence, P_2 preferably comprises a sequence which (1) binds a transition metal, (2) is hydrophobic to enhance the ability of the peptide to penetrate cell membranes and/or reach target tissues (*e.g.*, to be able to cross the blood brain barrier), or (3) otherwise stabilizes or enhances the performance of the peptide. P_2 together with P_1 may also be the N-terminal sequence of a protein having an N-terminal metal-binding site with high affinity for copper and nickel, such as human, rat or bovine serum albumin. In the case where $n = 100$, the peptide would have the sequence of approximately domain 1 of these albumins.

The sequences of many peptides which comprise a binding site for transition metal ions are known. See, *e.g.*, U.S. Patents Nos. 4,022,888, 4,461,724, 4,665,054, 4,760,051, 4,767,753, 4,810,693, 4,877,770, 5,023,237, 5,059,588, 5,102,990, 5,118,665, 5,120,831, 5,135,913, 5,145,838, 5,164,367, 5,591,711, 5,177,061, 5,214,032, 5,252,559, 5,348,943, 5,443,816, 5,538,945, 5,550,183, 5,591,711, 5,690,905, 5,759,515, 5,861,139, 5,891,418, 5,928,955, and 6,017,888, PCT applications WO 94/26295, WO 99/57262 and WO

99/67284, European Patent application 327263, Lappin et al., *Inorg. Chem.*, **17**, 1630-34 (1978), Bossu et al., *Inorg. Chem.*, **17**, 1634-40 (1978), Chakrabarti, *Protein Eng.*, **4**, 57-63 (1990), Adman, *Advances In Protein Chemistry*, **42**, 145-97 (1991), Cotelle et al., *J. Inorg. Biochem.*, **46**, 7-15 (1992), Canters et al., *FEBS*, **325**, 39-48 (1993), Regan, *Annu. Rev. Biophys. Biomol. Struct.*, **22**, 257-281 (1993), Ueda et al., *J. Inorg. Biochem.*, **55**, 123-30 (1994), Ueda et al., *Free Radical Biol. Med.*, **18**, 929-33 (1995), Regan, *TIBS*, **20**, 280-85 (1995), Ueda et al., *Chem. Pharm. Bull.*, **43**, 359-61 (1995), Bal et al., *Chem. Res. Toxicol.*, **10**, 906-914 (1997), Bal et al., *Chem. Res. Toxicol.*, **10**, 915-21 (1997), Koch et al., *Chem. Biol.*, **4**, 549-60 (1997), Kowalik-Jankowska et al., *J. Inorg. Biochem.*, **66**, 193-96 (1997), Harford and Sarkar, *Acc. Chem. Res.*, **30**, 123-130 (1997), Prince et al., *TIBS*, **23**, 197-98 (1998), Mlynarz, et al., *Speciation 98: Abstracts*, <http://www.jate.u-szeged.hu/~spec98/abstr/mlynar.html>, and Aitken, *Molec. Biotechnol.*, **12**, 241-53 (1999), Whittall et al., *Protein Science*, **9**, 332-343 (2000). P₂ may comprise the sequence of one or more of the metal-binding sites of these peptides.

When P₂ comprises a metal-binding site, it preferably has a sequence which includes a short spacer sequence between P₁ and the metal binding site of P₂, so that the metal-binding sites of P₁ and P₂ may potentially cooperatively bind metal ions (similar to a 2:1 peptide:metal complex; see Example 10). Preferably, the spacer sequence is composed of 1-5, preferably 1-3, neutral amino acids. Thus, the spacer sequence may be Gly, Gly Gly, Gly Ala Gly, Pro, Gly Pro Gly, etc.

In particular, when P₂ comprises a metal-binding site, it preferably comprises one of the following sequences: (Xaa₄)_m Xaa₃ His Xaa₂ Xaa₅, (Xaa₄)_m His Xaa₂ Xaa₅, (Xaa₄)_m Xaa₅ Xaa₂ His Xaa₃, or (Xaa₄)_m Xaa₅ Xaa₂ His. Xaa₂, Xaa₃ and Xaa₄ are defined above, and m is 0-5, preferably 1-3. The Xaa₄ amino acids form a short spacer sequence between P₁ and the metal binding site of P₂ so that the metal-binding sites of P₁ and P₂ may cooperatively bind metal ions, and Xaa₄ is preferably a neutral amino acid (see the previous paragraph). Xaa₅ is an amino acid having a free side-chain -NH₂, preferably Orn or Lys, more preferably Orn.

See Harford and Sarkar, *Acc. Chem. Res.*, **30**, 123-130 (1997) (the free side-chain -NH₂ of Orn within a peptide sequence has been reported to successfully substitute for the free N-terminal -NH₂ of the ATCUN motif). Thus, for instance, P₁ - P₂ could be Asp Ala His Gly Gly His Ala Orn [SEQ ID NO:2].

5 The amino acids of the peptide may be L-amino acids, D-amino acids, or a combination thereof. Preferably, at least one of the amino acids of P₁ is a D-amino acid (preferably Xaa₁ and/or His), except for β-Ala. Most preferably, all of the amino acids of P₁, other than β-Ala, are D-amino acids. Also, preferably about 50% of the amino acids of P₂ are D-amino acids, and most preferably all of the amino acids of P₂ are D-amino acids. D-amino acids are preferred because peptides containing D-amino acids are resistant to proteolytic enzymes, such as those that would be encountered upon administration of the peptide to an animal (including humans) or would be present in an excised organ perfused with a solution containing the peptide. Also, the use of D-amino acids would not alter the ability of the peptide to bind metal ions, including the ability of the peptide to bind copper with high affinity.

10 The peptides of the invention may be made by methods well known in the art. For instance, the peptides, whether containing L-amino acids, D-amino acids, or a combination of L- and D-amino acids, may be synthesized by standard solid-phase peptide synthesis methods. Suitable techniques are well known in the art, and include those described in Merrifield, in Chem. Polypeptides, pp. 335-61 (Katsoyannis and Panayotis eds. 1973); Merrifield, J. Am. Chem. Soc., **85**, 2149 (1963); Davis et al., Biochem. Int'l, **10**, 394-414 (1985); Stewart and Young, Solid Phase Peptide Synthesis (1969); U.S. Patents Nos. 3, 941,763 and 5,786,335; Finn et al., in The Proteins, 3rd ed., vol. 2, pp. 105-253 (1976); and Erickson et al. in The Proteins, 3rd ed., vol. 2, pp. 257-527 (1976). See also, Polish Patent 315474 (synthesis of HMS-containing peptides). Alternatively, the peptides may be synthesized by recombinant DNA techniques if they contain only L-amino acids. Recombinant DNA methods and suitable host cells, vectors and other reagents for use therein, are well known in the art. See, e.g., Maniatis et al., *Molecular Cloning: A*

Laboratory Manual, Cold Spring Harbor, NY (1982), Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, NY (1989).

The invention further comprises derivatives of the peptide $P_1 - P_2$, whether composed of L-amino acids, D-amino acids, or a combination of L- and D-amino acids, which are more resistant to proteolytic enzymes, more lipid soluble (to allow the peptides to more readily penetrate cell membranes and/or reach target organs, such as the brain), or both. As illustrated in Figure 1A, P_1 can be modified in the regions indicated by the arrows without altering the metal binding function of P_1 . In particular, P_1 can be substituted at carbons 1 or 2 with R_1 , and the terminal -COOH of P_1 can be substituted with protecting group R_2 (Figures 1B-D). P_2 can be modified in ways similar to those described for P_1 to make P_2 more resistant to proteolytic enzymes, more lipid soluble, or both.

R_1 can be a straight-chain or branched-chain alkyl containing from 1 to 16 carbon atoms, and the term "alkyl" includes the R and S isomers. R_1 can also be an aryl or heteroaryl containing 1 or 2 rings. The term "aryl" means a compound containing at least one aromatic ring (e.g., phenyl, naphthyl, and diphenyl). The term "heteroaryl" means an aryl wherein at least one of the rings contains one or more atoms of S, N or O. These substitutions do not substantially decrease the ability of P_1 to bind metal ions. In particular, the ability of P_1 to bind copper with high affinity is not decreased by these substitutions. For instance, some of the substituents, such as a n-butyl attached to carbon 2 (see Figure 1C, R_1 is n-butyl) should increase the affinity of the peptide for metal ions, such as copper, due to the inductive effect of the alkyl group. Substitution of carbon 2 (Figure 1C) with an aryl, heteroaryl, or a long chain alkyl (about 6-16 carbon atoms) should enhance transport of the peptide across lipid membranes.

As noted above, methods of synthesizing peptides by solid phase synthesis are well known. These methods can be modified to prepare the derivatives shown in Figures 1B-C. For example, the derivative of P_1 illustrated in Figure 1C, wherein R_1 is octyl, can be prepared as illustrated in Figure 2A. In Figure 2A, the elliptical element represents the polymer resin and R_p is a standard carboxyl protecting group. As illustrated in Figure 2A, octanoic acid

(freshly distilled) is treated with dry bromine followed by phosphorus trichloride. The mixture is heated to about 100°C and kept at that temperature for 4 hours. α -Bromooctanoic acid is obtained as a colorless liquid upon distillation. Amination of the bromoacid is achieved by allowing the acid and an ammonia solution to stand at 40-50° C for 30 hours. The octyl derivative of the amino acid is obtained by removing ammonium bromide with methanol washes. Classical resolution methods give the desired optically-pure D-form. Other derivatives wherein R_1 is an alkyl, aryl or heteroaryl can be prepared in the manner illustrated in Figure 2A.

In addition, the derivative of P_1 illustrated in Figure 1B, wherein R_1 is phenyl, can be prepared as illustrated in Figure 2B. In Figure 2B, Polymer is the resin, t-Bu is t-butyl, and Bz is benzyl. Other derivatives wherein R_1 is an alkyl, aryl or heteroaryl can be prepared in the manner illustrated in Figure 2B.

R_2 can be $-NH_2$, $-NHR_1$, $-N(R_1)_2$, $-OR_1$, or R_1 (see Figure 1D), wherein R_1 is defined above. These derivatives can be prepared as the last step of a solid-phase peptide synthesis before the peptide is removed from the resin by methods well known in the art. Substitutions with R_2 do not substantially decrease the ability of P_1 to bind metal ions.

In addition, P_1 and P_2 can be substituted with non-peptide functional groups that bind metal ions. These metal-binding functional groups can be attached to one or more pendent groups of the peptide, and the resulting peptide derivatives will possess one or more sites that are capable of binding metal ions, in addition to the binding site provided by P_1 and, optionally, the binding site provided by P_2 . As a consequence, the ability of such peptide derivatives to bind metal ions is improved as compared to the corresponding unmodified peptide. For instance, the peptide derivative can bind two of the same type of metal ion instead of one (*e.g.*, two Cu(II)), the peptide derivative can bind two different metal ions instead of one type of metal ion (*e.g.*, one Cu(II) and one Fe(III)), or the peptide derivative can bind one metal ion better (*e.g.*, with greater affinity) than the corresponding unmodified peptide.

Metal-binding functional groups include polyamines (*e.g.*, diamines, triamines, etc.). Suitable diamines include 1,2-alkyldiamines, preferably alkyl diamines wherein the alkyl contains 2-10 carbon atoms (*e.g.*, $\text{H}_2\text{N} - (\text{CH}_2)_n - \text{NH}_2$, wherein $n = 2-10$). Suitable diamines also include 1,2-aryldiamines, preferably benzene diamines (*e.g.*, 1,2-diaminobenzene). Suitable diamines further include 1,2-cyclic alkane diamines. "Cyclic alkanes" are compounds containing 1-3 rings, each containing 5-7 carbon atoms. Preferably the cyclic alkane diamine is 1,2-diaminocyclohexane (cyclohexane diamine).

A particularly preferred diamine is 1,2-diaminocyclohexane (Figures 3A-B). Previous studies carried out by Rao & P. Williams (*J. Chromatography A*, **693**, 633 (1995)) have shown that a cyclohexane diamine derivative (Figure 3A, where PYR is pyridine) binds to a variety of metal ions. The resulting metal chelator has been successfully used to resolve amino acids and peptides, showing that the molecule has a very high affinity for α -amino acids, forming a very stable coordination complex, which is unique in many respects. 1,2-Diaminocyclohexane possesses a reactive amino functional group to which a peptide of the invention can be attached. See Figure 3B, where M is a metal ion and at least one R_4 is -alkyl-CO-peptide, -aryl-CO-peptide, -aryl-alkyl-CO-peptide, or -alkyl-aryl-CO-peptide (see also Figures 3C-D). The other R_4 may be the same or may be -alkyl-COOH, -aryl-COOH, -aryl-alkyl-COOH, or alkyl-aryl-COOH. Derivatives of the type shown in Figure 3B will have several metal-binding sites and can, therefore, be expected to bind metal ions more readily than the unsubstituted peptide. Further, due to the presence of the cyclohexane functionality, the compound will possess lipid-like characteristic which will aid its transport across lipid membranes.

Cyclohexane diamine derivatives of the peptides of the invention can be prepared by two distinct routes. The first involves initial condensation with an aldehyde followed by reduction (see Figure 3C; in Figure 3C Bz is benzyl). A number of aldehydes (alkyl and aryl) react readily with cyclohexane diamine at room temperature, forming an oxime. The oxime can be reduced with sodium borohydride under anaerobic conditions to give the diacid derivative. The carboxyl moieties are then reacted with the free amino groups present in

carboxy-protected P_1 to give the cyclohexane diamine derivative of the peptide. The second route is a direct alkylation process which is illustrated in Figure 3D. For example, cyclohexane diamine is treated with bromoacetic acid to give the diacetic acid derivative. The carboxyl moieties are then reacted with the free amino groups present in carboxy-protected P_1 to give the derivative. In Figure 3D, R_5 is H or another peptide. When R_5 is H, the derivative can be further reacted to produce typical carboxylic acid derivatives, such as esters, by methods well known in the art. Metal binding experiments have indicated that the presence or absence of this group does not have a bearing on the metal binding capacity of the whole molecule. However, these groups would either make the molecule hydrophobic or hydrophilic, depending upon the substituent, and this may, in turn, have an effect on delivery of the molecule across membranes or to target tissues. These two synthetic routes will work for the synthesis of diamine peptide derivatives using the other diamines described above.

Additional suitable polyamines and polyamine derivatives and methods of attaching them to peptides are described in U.S. Patents Nos. 5,101,041 and 5,650,134, the complete disclosures of which are incorporated herein by reference. Other polyamine chelators suitable for attachment to peptides are known. See, *e.g.*, U.S. Patents Nos. 5,422,096, 5,527,522, 5,628,982, 5,874,573, and 5,906,996 and PCT applications WO 97/44313, WO 97/49409, and WO 99/39706.

It is well known that vicinal diacids bind to metal ions, and the affinity for copper is particularly high. It is therefore envisaged that a peptide having a vicinal diacid functional group will be extremely effective in metal binding. Suitable vicinal diacids include any 1,2-alkyldiacid, such as diacetic acid (succinic acid), and any 1,2-aryldiacid.

The amino groups of the peptide can be reacted with diacetic acid to produce a diacid derivative (see Figure 4). This can be conveniently accomplished by reacting the amino groups of the resin-bound peptide with a halogenated acetic acid (*e.g.*, bromoacetic acid or chloroacetic acid) or a halogenated acetic acid derivative (*e.g.*, benzyloxy ester). Solid phase synthetic procedures enable removal of unreacted materials by washing with solvent. The

final product is released from the resin by hydrolytic cleavage. Other diacid derivatives of the peptides of the invention can be made in the same manner.

Polyaminopolycarboxylic acids are known to bind metals, such as copper and iron. Suitable polyaminopolycarboxylic acids for making derivatives of the peptides of the invention and methods of attaching them to peptides are described in U.S. Patents Nos. 5,807,535 and 5,650,134, and PCT application WO 93/23425, the complete disclosures of which are incorporated herein by reference. See also, U.S. Patent No. 5,739,395.

Vicinal polyhydroxyl derivatives are also included in the invention. Suitable vicinal polyhydroxyls include monosaccharides and polysaccharides (*i.e.*, disaccharide, trisaccharide, etc.). Presently preferred are monosaccharides. See Figure 7. The monosaccharides fall into two major categories - furanoses and pyranoses. One of the prime examples of a furanose ring system is glucose. The hydroxyl groups of glucose can be protected as benzyl or labile *t*-butoxy functional groups, while leaving the aldehyde free to react with an amine group (*e.g.*, that of lysine) of the tetrapeptide. Mild reduction/hydrolysis produces the monosaccharide peptide derivative. Other monosaccharide peptide derivatives can be prepared in this manner.

Bispyridylethylamine derivatives are known to form strong complexes with divalent metal ions. When attached to the peptide, this functional group would provide additional chelating sites for metal ions, including copper. The bispyridylethyl derivative of the tetrapeptide Asp Ala His Lys [SEQ ID NO:1] is shown in Figure 5. It is anticipated that the metal-binding capacity of this tetrapeptide derivative will be increased by at least three-fold as compared to the underivatized peptide. The preparation of this bispyridylethylamine derivative shares some similarities with the synthesis of diacid derivatives. The two amino groups of the tetrapeptide (one at Asp and the other at Lys) are reacted with 2-bromoethylpyridine to give the tetra-substituted peptide derivative. The reaction is accomplished by reacting the resin-bound tetrapeptide with the bromoethylpyridine, followed by cleavage of the product from the resin.

Phenanthroline is another heterocyclic compound capable of binding divalent metal ions. Phenanthroline derivatives of the peptides can be synthesized in the same manner as for the bispyridylethylamine derivatives.

5 Porphyrins are a group of compounds found in all living matter and contain a tetrapyrrolic macrocycle capable of binding to metals. Heme, chlorophyll and corrins are prime examples of this class of compounds containing iron, magnesium and cobalt, respectively. Mesoporphyrin IX (Figure 6A-B, where M is a metal ion) is derived from heme and has been observed to possess specific affinity for copper. Addition of this structure to a peptide of the invention would produce a porphyrin-peptide derivative possessing several
10 sites for binding of copper (see Figure 6C). In addition to their roles in metal binding, the imidazole residues at positions 3 and 3' of the tetrapeptide shown in Figure 6C may provide a binding site for metals other than copper, thereby stabilizing the porphyrin-metal complex. In particular, cyanocobalamine (vitamin B-12) contains cobalt as the metal in the porphyrin nucleus, and the complex is stabilized by the imidazole groups. On the basis of this analogy
15 it is anticipated that the porphyrin-tetrapeptide derivative would bind cobalt (or other metals) at normal physiological conditions in the porphyrin nucleus and that the complex would be stabilized by the His imidazole groups.

To prepare the porphyrin-peptide derivative shown in Figure 6C, the carboxyl groups of mesoporphyrin IX can be activated and coupled with the amino groups of the peptide employing standard solid-phase peptide synthesis. Typically, the free amino group of the lysine residue of the resin-bound peptide can be coupled with carboxy activated porphyrin
20 nucleus. The condensation product can be cleaved off the resin using standard methods. This method can be used to synthesize other porphyrin derivatives of peptides of the invention.

Other suitable porphyrins and macrocyclic chelators and methods of attaching them
25 to peptides are described in U.S. Patents Nos. 5,994,339 and 5,087,696, the complete disclosures of which are incorporated herein by reference. Other porphyrins and macrocyclic chelators that could be attached to peptides are known. See, *e.g.*, U. S. Patents Nos.

5,422,096, 5,527,522, 5,628,982, 5,637,311, 5,874,573, and 6,004,953, PCT applications WO 97/44313 and WO 99/39706.

A variety of additional metal chelators and methods of attaching them to proteins are described in U.S. Patent No. 5,683,907, the complete disclosure of which is incorporated herein by reference.


Dithiocarbamates are known to bind metals, including iron. Suitable dithiocarbamates for making derivatives of the peptides of the invention are described in U.S. Patents Nos. 5,380,747 and 5,922,761.

Hydroxypyridones are also known to be iron chelators. Suitable hydroxypyridones for making derivatives of the peptides of the invention are described in U.S. Patents Nos. 4,912,118 and 5,104,865 and PCT application WO 98/54138.

Additional non-peptide metal chelators are known in the art or will be developed. Methods of attaching chemical compounds to proteins and peptides are well known in the art, and attaching non-peptide metal chelators to the peptides of the invention is within the skill in the art. See, *e.g.*, those patents cited above describing such attachment methods.

As can be appreciated, the non-peptide, metal-binding functional groups could be attached to another metal-binding peptide (MBP) in the same manner as they are to peptide P₁-P₂. The resulting peptide derivatives would contain one or more metal-binding functional groups in addition to the metal-binding site of MBP. Preferably, MBP contains from 2-10, more preferably 3-5, amino acids. Preferably MBP contains one or more D-amino acids; most preferably all of the amino acids of MBP are D-amino acids. As described above, the sequences of many metal-binding peptides are known. These peptides and peptides comprising the metal-binding sites of these peptides can be prepared in the same ways as described above for peptide P₁-P₂. Derivatives of these peptides having one or more metal-binding functional group attached to the peptide can be prepared in the same ways as described above for derivatives of peptide P₁-P₂.

The invention also provides metal-binding peptide dimers of the formula:



P_3 is any peptide capable of binding a metal ion, and each P_3 may be the same or different. Each P_3 preferably contains 2-10, more preferably 3-5, amino acids. As described above, metal-binding peptides are known, and each P_3 may comprise the sequence of one or more of the metal-binding sites of these peptides. Although each P_3 may be substituted as described above for P_1 and P_2 , including with a non-peptide, metal-binding functional group, both P_3 peptides are preferably unsubstituted. P_3 may also comprise any amino acid sequence substituted with a non-peptide, metal-binding functional group as described above to provide the metal-binding capability of P_3 . Preferably, each P_3 is an unsubstituted metal-binding peptide (*i.e.*, an unsubstituted peptide comprising a peptide sequence which binds metal ions). Most preferably, one or both of the P_3 groups is P_1 (*i.e.*, the dimers have the sequence P_3 - S - P_1 , P_1 - S - P_3 or, most preferably, P_1 - S - P_1). P_1 is defined above.

L is a linker which is attached to the C-terminal amino acid of each P_3 . L may be any chemical group which can connect the two P_3 peptides through their C-terminal amino acids which is physiologically-acceptable. By "physiologically-acceptable" is meant that a peptide dimer containing the linker L is not toxic to an animal (including a human) or an organ to which the peptide dimer is administered as a result of the inclusion of the linker L in the peptide dimer. Preferably, L links the two P_3 groups so that they can cooperatively bind metal ions (similar to a 2:1 peptide:metal complex; see Example 10). L is also preferably neutral. Most preferably, L is a straight-chain or branched-chain alkane or alkene residue containing from 1-18, preferably from 2-8, carbon atoms (*e.g.*, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2\text{CH}_2-$, $-\text{CH}_2\text{CH}_2(\text{CH}_3)\text{CH}_2-$, $-\text{CHCH}-$, etc.) or a cyclic alkane or alkene residue containing from 3-8, preferably from 5-6, carbon atoms (see Figure 19A, compound D_1), preferably attached to a P_3 by means of an amide linkage. Such linkers are particularly preferred because they impart hydrophobicity to the peptide dimers. In another preferred embodiment, L is a nitrogen-containing heterocyclic alkane residue (see Figure 19A, compounds D_2 , D_3 and D_4), preferably a piperazide (see Figure 19A, compound D_2). In another preferred embodiment L is a glyceryl ester (see Figure 19A, compound D_5 ; in formula D_5 , R is an alkyl or aryl containing, preferably containing 1-6 carbon atoms). These preferred linkers L will allow the two peptides P_3 to bind

metal ions cooperatively and are biocompatible, and the peptide dimers can be made easily and in large quantities. By “biocompatible” is meant that a peptide dimer containing the linker L does not produce any undesirable side-effects due to the linker L in an animal (including a human) or an organ to which the peptide dimer is administered.

5 Methods of synthesizing the peptide dimers are illustrated in Figures 19B-D. In general, the C-terminal amino acids (protected by methods and protecting groups well known in the art) of the two P₃ groups are attached to L, and the resulting amino acid dimers used in standard peptide synthetic methods to make the peptide dimers.

10 For instance, a peptide dimer, where each peptide has the sequence Asp Ala His Lys, can be synthesized by coupling protected lysines to a free diamine functional group, either as an acid chloride or by using standard coupling agents used in peptide synthesis (see Figures 19B-C). Many suitable diamines are available commercially or suitable diamines can be readily synthesized by methods known in the art.

15 For instance, the lysine dimer **2** (Figure 19B) can be prepared as follows. To a stirred solution of 9-fluorenylmethyloxycarbonyl (Fmoc)- and t-benzyloxycarbonyl(Boc)- protected D-Lys (Fmoc-D-Lys(Boc)-OH) (20 mmole) in dry dimethylformamide (DMF; 100 mL; dry argon flushed) are added butane-1,4-diamine **1** and 2-(1H-benzotriazole-1-yl)-1,2,3,3-tetramethyluroniumtetrafluoroborate (TBTU; 0.5 mmole). The solution is stirred for 36 hours at room temperature. The bis-protected lysine **2** is isolated by flash chromatography over silica and elution with mixtures of ethyl acetate/methanol. The peptide dimer **3** is then prepared from the protected lysine dimer **2** employing classical peptide synthesis methodology (see Figure 19B).

20 Another peptide dimer, where each peptide has the sequence Asp Ala His Lys, can be synthesized as follows. First, a different protected lysine dimer **4** is synthesized by acylating the two amino centers of a piperazine **5** (see Figure 19C; see also Chambrier et al., *Proc. Natl. Acad. Sci.*, **96**, 10824-10829 (1999)). Then, the remainder of the amino acid residues are added employing standard peptide synthesis methodology to give the peptide dimer **6** (see Figure 19C).

Peptide dimers, where each peptide has the sequence Asp Ala His Lys and where L is a glyceryl ester, can be synthesized as follows. The 3-substituted propane-1,2-diols of formula 7 in Figure 19D, wherein R is an alkyl or aryl, are commercially available. A lysine diester 8, wherein R is methyl, can be prepared as follows (see Figure 19D). To a stirred solution of Fmoc-D-Lys(Boc)-OH (20 mmole) in dry toluene (100 mL; dry argon flushed) is added 3-methoxypropane-1,2-diol (200 mmole) and imidazole (15 mmole). The solution is stirred for 36 hours at room temperature. The solvent is removed in vacuo, and the residue is dissolved in ethyl acetate. This solution is washed with citric acid solution (2%), water, 0.5 N NaHCO₃ solution, and again with water; then the organic layer is dried over magnesium sulphate (removal of the solvent gives a pale yellow residue). The bis-protected lysine 8 is isolated by flash chromatography over silica and elution with mixtures of ethyl acetate/methanol. The peptide dimer 9 is then prepared from the protected lysine dimer 8 employing classical peptide synthesis methodology (see Figure 19D).

The physiologically-acceptable salts of the metal-binding compounds are also included in the invention. Physiologically-acceptable salts include conventional non-toxic salts, such as salts derived from inorganic acids (such as hydrochloric, hydrobromic, sulfuric, phosphoric, nitric, and the like), organic acids (such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, glutamic, benzoic, salicylic, and the like) or bases (such as the hydroxide, carbonate or bicarbonate of a pharmaceutically-acceptable metal cation). The salts are prepared in a conventional manner, *e.g.*, by neutralizing the free base form of the compound with an acid.

A metal-binding compound of the invention can be used to reduce the damage done by ROS or to reduce excess metal ion concentrations in an animal. To do so, a metal-binding compound of the invention is administered to the animal. Preferably, the animal is a mammal, such as a rabbit, goat, dog, cat, horse or human. Effective dosage forms, modes of administration and dosage amounts for the various compounds of the invention may be determined empirically, and making such determinations is within the skill of the art. It has been found that an effective dosage is from about 2 to about 200 mg/kg, preferably from about

10 to about 40 mg/kg, most preferably about 20 mg/kg. However, it is understood by those skilled in the art that the dosage amount will vary with the particular metal-binding compound employed, the disease or condition to be treated, the severity of the disease or condition, the route(s) of administration, the rate of excretion of the compound, the duration of the treatment, the identify of any other drugs being administered to the animal, the age, size and species of the animal, and like factors known in the medical and veterinary arts. In general, a suitable daily dose of a compound of the present invention will be that amount of the compound which is the lowest dose effective to produce a therapeutic effect. However, the daily dosage will be determined by an attending physician or veterinarian within the scope of sound medical judgment. If desired, the effective daily dose may be administered as two, three, four, five, six or more sub-doses, administered separately at appropriate intervals throughout the day. Administration of the compound should be continued until an acceptable response is achieved.

The compounds of the present invention may be administered to an animal patient for therapy by any suitable route of administration, including orally, nasally, rectally, vaginally, parenterally (*e.g.*, intravenously, intraspinally, intraperitoneally, subcutaneously, or intramuscularly), intracisternally, transdermally, intracranially, intracerebrally, and topically (including buccally and sublingually). The preferred routes of administration are orally and intravenously.

While it is possible for a metal-binding compound of the present invention to be administered alone, it is preferable to administer the compound as a pharmaceutical formulation (composition). The pharmaceutical compositions of the invention comprise a metal-binding compound or compounds of the invention as an active ingredient in admixture with one or more pharmaceutically-acceptable carriers and, optionally, with one or more other compounds, drugs or other materials. Each carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not injurious to the animal. Pharmaceutically-acceptable carriers are well known in the art. Regardless of the route of administration selected, the compounds of the present invention are formulated into

pharmaceutically-acceptable dosage forms by conventional methods known to those of skill in the art. See, e.g., *Remington's Pharmaceutical Sciences*.

Formulations of the invention suitable for oral administration may be in the form of capsules, cachets, pills, tablets, powders, granules or as a solution or a suspension in an aqueous or non-aqueous liquid, or an oil-in-water or water-in-oil liquid emulsions, or as an elixir or syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia), and the like, each containing a predetermined amount of a compound of the present invention as an active ingredient. A compound of the present invention may also be administered as bolus, electuary or paste.

In solid dosage forms of the invention for oral administration (capsules, tablets, pills, dragees, powders, granules and the like), the active ingredient is mixed with one or more pharmaceutically acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; (3) humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin; (6) absorption accelerators, such as quaternary ammonium compounds; (7) wetting agents, such as, for example, cetyl alcohol and glycerol monostearate; (8) absorbents, such as kaolin and bentonite clay; (9) lubricants, such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof; and (10) coloring agents. In the case of capsules, tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid compositions of a similar type may be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

A tablet may be made by compression or molding optionally with one or more accessory ingredients. Compressed tablets may be prepared using binder (for example, gelatin or hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (for

example, sodium starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

5 The tablets, and other solid dosage forms of the pharmaceutical compositions of the present invention, such as dragees, capsules, pills and granules, may optionally be scored or prepared with coatings and shells, such as enteric coatings and other coatings well known in the pharmaceutical-formulating art. They may also be formulated so as to provide slow or controlled release of the active ingredient therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile, other polymer matrices, liposomes and/or microspheres. They may be sterilized by, for example, filtration through a bacteria-retaining filter. These compositions may also optionally contain opacifying agents and may be of a composition that they release the active ingredient only, or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. The active ingredient can also be in microencapsulated form.

10 Liquid dosage forms for oral administration of the compounds of the invention include pharmaceutically-acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredient, the liquid dosage forms may contain inert diluents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

15 Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring, perfuming and preservative agents.

20 Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters,

microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

Formulations of the pharmaceutical compositions of the invention for rectal or vaginal administration may be presented as a suppository, which may be prepared by mixing one or more compounds of the invention with one or more suitable nonirritating excipients or carriers comprising, for example, cocoa butter, polyethylene glycol, a suppository wax or salicylate, and which is solid at room temperature, but liquid at body temperature and, therefore, will melt in the rectum or vaginal cavity and release the active compound. Formulations of the present invention which are suitable for vaginal administration also include pessaries, tampons, creams, gels, pastes, foams or spray formulations containing such carriers as are known in the art to be appropriate.

Dosage forms for the topical or transdermal administration of a compound of this invention include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches, drops and inhalants. The active compound may be mixed under sterile conditions with a pharmaceutically-acceptable carrier, and with any buffers, or propellants which may be required.

The ointments, pastes, creams and gels may contain, in addition to an active compound of this invention, excipients, such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

Powders and sprays can contain, in addition to a compound of this invention, excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder or mixtures of these substances. Sprays can additionally contain customary propellants such as chlorofluorohydrocarbons and volatile unsubstituted hydrocarbons, such as butane and propane.

Transdermal patches have the added advantage of providing controlled delivery of a compound of the invention to the body. Such dosage forms can be made by dissolving, dispersing or otherwise incorporating a compound of the invention in a proper medium, such

as an elastomeric matrix material. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate of such flux can be controlled by either providing a rate-controlling membrane or dispersing the compound in a polymer matrix or gel.

Pharmaceutical compositions of this invention suitable for parenteral administrations comprise one or more compounds of the invention in combination with one or more pharmaceutically-acceptable sterile isotonic aqueous or non-aqueous solutions, dispersions, suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain antioxidants, buffers, solutes which render the formulation isotonic with the blood of the intended recipient or suspending or thickening agents.

Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

These compositions may also contain adjuvants such as wetting agents, emulsifying agents and dispersing agents. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like in the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

In some cases, in order to prolong the effect of a drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material having poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively,

delayed absorption of a parenterally-administered drug is accomplished by dissolving or suspending the drug in an oil vehicle.

Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending on the ratio of drug to polymer, and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissue. The injectable materials can be sterilized for example, by filtration through a bacterial-retaining filter.

The formulations may be presented in unit-dose or multi-dose sealed containers, for example, ampules and vials, and may be stored in a lyophilized condition requiring only the addition of the sterile liquid carrier, for example water for injection, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the type described above.

As noted above, ROS have been reported to play a major role in a variety of diseases and conditions. See Manso, *Rev. Port. Cardiol.*, **11**, 997-999 (1992); Florence, *Aust. N Z J. Ophthalmol.*, **23**, 3-7 (1992); Stohs, *J. Basic Clin. Physiol. Pharmacol.*, **6**, 205-228 (1995); Knight, *Ann. Clin. Lab. Sci.*, **25**, 111-121 (1995); Kerr et al., *Heart & Lung*, **25**, 200-209 (1996). Diseases involving or caused by excess metal ions are also known. The metal-binding compounds of the invention can be used to treat any of these diseases and conditions and other diseases and conditions in which ROS or transition metals play a role. Specific diseases and conditions treatable with the metal-binding compounds of the invention include adult respiratory distress syndrome, aging, AIDS, atherosclerosis (hypertension, senility and impotence), arthritis, asthma, autoimmune diseases, cancer (*e.g.*, kidney, liver, colon, and brain), carcinogenesis, cellular damage caused by ionizing radiation (*e.g.*, radiation of tumors), chronic granulomatous disease, cirrhosis, Crohn's disease, diabetes (diabetic retinopathy, renal disease, impotence and peripheral vascular disease), eye diseases (*e.g.*, cataracts, central artery occlusion, benign monoclonal gammopathy, and macular degeneration), emphysema,

inflammation, ischemia, neoplastic diseases, neurological trauma, neurodegenerative diseases (e.g., Alzheimer's disease, amyotrophic lateral sclerosis, Parkinson's disease, multiple sclerosis, and senile dementia), pancreatitis, peripheral vascular disease, pulmonary embolism, renal disease (dialysis patients), reperfusion, shock, tissue damage occurring upon administration of chemotherapeutics, tissue damage after surgery (e.g., transplantation surgery, open heart surgery, and any surgery where the blood supply to a tissue is cut off, and surgical ischemia of the limbs (tourniquet injury)), toxic reactions (e.g. herbicide poisoning, transition metal (copper, cobalt, and nickel) poisoning, carbon monoxide poisoning, and antibiotic toxicity), traumatic crush injury, and Wilson's disease (congenital high levels of copper). Specific ischemic conditions and diseases treatable with the metal-binding compounds of the invention include:

Central nervous system ischemia -

Brain ischemia after surgery

Hyperthermia brain injury

Perinatal hypoxia-induced ischemia ("cerebral palsy")

Spinal cord injury

Stroke (thrombotic, embolic or hemorrhagic cerebrovascular accident)

Transient ischemic attack

Traumatic brain injury

Cardiac ischemia -

Acute myocardial infarction

Angina pectoris

Arrhythmias

Cardiac ischemia after surgery

Congestive heart failure

Myocardial "stunning"

Ischemic bowel disease

Placental ischemia

Pulmonary embolism

Surgery where the blood supply to a tissue or organ is cut off -

Angioplasty

Cardiac bypass surgery

5 Transplantation surgery (both the donor organ and the recipient of the organ)

Surgical ischemia of the limbs (tourniquet injury).

A compound of the invention is preferably administered prophylactically. For instance, a compound of the invention is preferably administered prior to and/or simultaneously with
 10 reperfusion of an ischemic tissue or organ (*e.g.*, prior to and/or simultaneously with angioplasty or treatment with clot dissolving drugs, such as tissue plasminogen activators). Of course administration of a compound of the invention should be continued for a period of time after reperfusion has been achieved. Similarly, a compound of the invention should be administered prior to and/or during surgery (*e.g.*, open-heart surgery or surgery to transplant
 15 an organ into an animal), and administration of the compound should be continued for a period of time after the surgery. As another example of prophylactic administration, a compound of the invention can be administered to a patient presenting symptoms of a serious condition (*e.g.*, cerebrovascular ischemia or cardiovascular ischemia) while the patient is tested to diagnose the condition. In this way, the patient will be protected during the time it takes to diagnose such conditions, and treatment with the metal-binding compounds of the invention may also
 20 prolong the time during which other therapies (*e.g.*, administration of tissue plasminogen activator for cerebrovascular ischemia) can be administered. As yet a further example, a compound of the invention can be administered at the time a patient is to undergo radiation therapy (*e.g.*, radiation for a tumor or prior to a bone marrow transplant).

25 A compound of the invention can also be used to treat patients who have suffered blunt trauma. In particular, a compound of the present invention may be very beneficial in treating patients suffering from multiple blunt trauma who have a low albumin level, since it has been found that a low albumin level is a predictor of mortality in such patients. More specifically,

34 patients suffering from multiple blunt trauma were studied. These patients were admitted to the intensive care unit of Swedish Hospital, Denver, CO in 1998. Two groups of patients were matched by a trauma surgeon by age, injury severity score (ISS), and type and area of injury without knowledge of the albumin levels of the patients. One group was composed of the patients who died, and the other group was composed of survivors. Following the match, the admission albumin levels were retrieved from the medical records by an independent observer, and the albumin levels of the two groups were compared. For the 17 survivors, the mean albumin level was 3.50 ± 1.00 g/dl. For the 17 patients who died, the mean albumin level was 2.52 ± 0.73 g/dl. The % variance was 28.6 and 28.9, respectively, and the p-value was 0.0026 (95% confidence interval 0.3462 - 0.4771).

The compounds of the invention may be given alone to reduce the damage done by ROS. Alternatively, the compounds of the invention can be given in combination with "free radical scavengers." "Free radical scavengers" include superoxide dismutase, catalase, glutathione peroxidase, ebselen, glutathione, cysteine, N-acetyl cysteine, penicillamine, allopurinol, oxypurinol, ascorbic acid, α -tocopherol, Trolox (water-soluble α -tocopherol), β -carotene, fatty-acid binding protein, fenozan, probucol, cyanidanol-3, dimercaptopropanol, indapamide, emoxipine, dimethyl sulfoxide, and others. See, e.g., Das et al., *Methods Enzymol.*, **233**, 601-610 (1994); Stohs, *J. Basic Clin. Physiol. Pharmacol.*, **6**, 205-228 (1995). The compounds of the invention can, of course, also be given along with standard therapies for a given conditions (e.g., insulin to treat diabetes).

The metal-binding compounds of the invention can also be used to reduce the damage done by ROS in a tissue or organ that has been removed from an animal. To do so, the tissue or organ is contacted with a solution containing an effective amount of a metal-binding compound of the invention. Many suitable solutions are known. See, e.g., Dunphy et al., *Am. J. Physiol.*, **276**, H1591-H1598 (1999); Suzer et al., *Pharmacol. Res.*, **37**, 97-101 (1998); Hisatomi et al., *Transplantation*, **52**, 754-755 (1991); U.S. Patent No. 5,710,172. Effective amounts of the peptide to include in such solutions can be determined empirically, and doing so is within the skill in the art. The harvested tissue or organ may subsequently be used for

transplantation into a recipient or for research purposes (*e.g.*, using a perfused liver to screen drugs).

5 The invention further provides a kit for reducing the damage done by ROS in a tissue or organ that has been removed from an animal. The kit is a packaged combination of one or more containers holding reagents and other items useful for preserving harvested organs. The kit comprises a container holding a metal-binding compound of the invention. Suitable containers include bottles, bags, vials, test tubes, syringes, and other containers known in the art. The kit may also contain other items which are known in the art and which may be desirable from a commercial and user standpoint, such as diluents, buffers, empty syringes, 10 tubing, gauze pads, disinfectant solution, etc.

11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642

EXAMPLES

EXAMPLE 1: Synthesis of Tetrapeptide Asp Ala His Lys [SEQ ID NO:1]

This example describes the synthesis of the tetrapeptide Asp Ala His Lys [SEQ ID NO:1] composed of all L-amino acids using standard solid-phase synthesis techniques. First, 9-fluorenylmethyloxycarbonyl (Fmoc)-protected Asp (v COO- ester; Tolsulfonyl) on Wang resin (0.6 mmole; Nova Biochem) was suspended in a solution of piperidine/dimethylformamide (DMF) (40% v/v; 3 ml) for 30 min with occasional agitation. At the end of this period, the solvent was drained, and the resin was washed sequentially with DMF and dichloromethane (DCM; 5 x 3 ml). A ninhydrin test was used to monitor the reaction. The resin was swollen with DMF (~1 ml). The C-protected t-benzyloxycarbonyl (Boc) ester of alanine in DMF was added, followed by a mixture of diisopropylamine (8 equivalent) and 2-(1H-benzotriazole-1-yl)-1,2,3,3-tetramethyluroniumtetrafluoroborate (TBTU-) (4 equivalents). The resin was shaken for about 24 hours, and the reaction was monitored by the ninhydrin test. At the end of this period, DMF was drained, and the resin was washed with DMF and DCM. The solution was drained, and the beads were washed with DCM (3 x 2 ml). The protecting group of the dipeptide-resin was removed, and the beads were suspended in DMF. Amino protected (benzyloxy) derivative of histidine (4 mmole) was added, followed by mixture of diisopropylamine (8 equivalent) and TBTU- (4 equivalent). The resin was shaken for about 24 hours, and the reaction monitored by ninhydrin test. At the end of this period, DMF was drained, and the resin was washed with DMF and DCM. The tripeptide-resin was briefly dried in a gentle stream of nitrogen and suspended in nitrogen-saturated DMF. Protected lysine was added, followed by a mixture of diisopropylamine (8 equivalent) and TBTU- (4 equivalent). The resin was shaken for about 24 hours, and the reaction monitored by the ninhydrin test. At the end of this period, DMF was drained and the resin was washed with DMF and DCM. The Boc protecting group was carefully removed to give the tetrapeptide bound to the resin, with a typical loading of 5 mmole/g. The resin bound tetrapeptide (0.25 gm; 5 mmolar) was treated with trifluoroacetic acid (TFA) and was shaken

for 24 hours. At the end of this period, the ninhydrin test gave a blue color, indicating the release of the tetrapeptide from the resin. In some circumstances, addition of 5%(V/V) of DMF to TFA accelerated the rate of release of the peptide from the resin. Removal of TFA at reduced pressure gave the tetrapeptide (all D) as TFA salt and was dried under vacuum at 5°C for 24 hours. The residue was a white powder and was characterized by spectrometric methods.

A number of enantiomers of the tetrapeptide can be prepared in this manner. For example, use of D-amino acids in the peptide synthesis forms the tetrapeptide containing all D-amino acids. Also, combinations of L-amino acids and D-amino acids can be used.

EXAMPLE 2: Preparation of Cyclohexanediamine

Derivative of Asp Ala His Lys [SEQ ID NO:1]

Trans-diaminocyclohexane was prepared by resolving cis/trans 1,2-diaminocyclohexane (Aldrich-Sigma) as the tartaric acid salt. The R-trans isomer melts at 75°C and the S-trans isomer melts between 43-45°C (Ph.D. Thesis, P.D. Newman, University College, Cardiff, U.K., 1994). The trans-diaminocyclohexane (10 gm) was then suspended in anhydrous toluene (30mL) and cooled to 5°C in an ice bath, and bromoacetic acid (8 gm) in toluene (25 mL) was added dropwise. At the end of the addition, the reaction temperature was raised to 30°C and kept at that temperature for a further 5 hours. Toluene was evaporated, and the R-trans 1,2-diaminocyclohexane diacetic acid was crystallized from hexane/toluene to give a white solid (yield 70%). The product was characterized by spectroscopic methods.

The resin-bound tetrapeptide prepared in Example 1 (20mg) was suspended in DMF (5 mL) and was treated with the R-trans 1,2-diaminocyclohexanediacytic acid (20 mg) followed by addition of a mixture of diisopropylamine (8 equivalent) and TBTU-(4 equivalent). The resin was shaken for about 24 h on a roller. Then, the resin was washed with DMF followed by DCM (5x3mL) and partially dried. Hydrolysis of the resin linkage was effected by treating the resin-bound reaction product with TFA (5mL; 5 hr). The resin was separated and washed with DCM. The washings were combined with TFA and concentrated under

vacuum. The residue (cyclohexanediamine tetrapeptide; formula given in Figure 3D where R₅ is H) was characterized by spectrometric analysis.

EXAMPLE 3: Preparation of Tetrapeptide Tetracetic Acid

The resin-bound tetrapeptide prepared in Example 1 (20 mg) was suspended in DMF (5 mL) and treated with excess (10-fold) chloroacetic acid. The resin was shaken at room temperature for 48 hours, followed by heating to 60°C for a further hour. DMF was removed by filtration, and the resin was washed with DMF followed by DCM (5x3mL). Partially dried resin was used without further treatment in the next stage. Hydrolysis of the resin linkage was effected by treating the resin-bound reaction product with TFA (5mL; 5 hr). The resin was separated and washed with DCM. The washings were combined with TFA and concentrated under vacuum (yield 30%). The product (formula given in Figure 4) was characterized by spectrometric methods.

EXAMPLE 4: Preparation of Mesoporphyrin IX Tetrapeptide

The resin-bound tetrapeptide prepared in Example 1 (20 mg) was suspended in DMF (5 mL) and treated with mesoporphyrin IX dicarboxylic acid (10 μmole; formula given in Figure 6A), followed by addition of a mixture of diisopropylamine (8 equivalent) and TBTU-(4 equivalent). The resin was shaken for about 24 hours on a roller kept in a dark chamber. The resin was washed with DMF followed by DCM (5x3mL) and partially dried. Hydrolysis of the resin linkage was effected by treating the resin-bound reaction product with TFA (5mL; 5 hr). The resin was separated and washed with DCM/TFA mixture (1:1.5mL). The washings were combined and concentrated under vacuum. The porphyrin tetrapeptide (formula given in Figure 6C) was purified by semi-preparative HPLC (yield 60%). The structure was confirmed by spectrometric methods.

This procedure can be used to synthesize other porphyrin-peptides, such as mesoporphyrin I and related molecules.

EXAMPLE 5: Preparation of Tetrabispiridylethyl Tetrapeptide

The resin-bound tetrapeptide prepared in Example 1 (20 mg) was suspended in DMF (5 mL) and treated with bromoethylpyridine (20 μ mole). This was followed by the addition of pyridine (0.5 mL). The resin was shaken for about 48 hours on a roller. The resin was washed with DMF, followed by DCM (5x3mL) to remove all of the unreacted monomers, and then dried under vacuum for 30 minutes. Hydrolysis of the resin linkage was effected by treating the resin-bound reaction product with TFA (5mL; 5 hr). The resin was separated and washed with DCM/TFA mixture (1:1.5mL). The washings were combined and concentrated under vacuum. The pyridylethyl tetrapeptide derivative (formula given in Figure 5) was purified by semi-preparative HPLC (yield 50 %). The structure was confirmed by spectrometric methods.

This procedure can be applied to other heterocycles, such as phenanthroline and related molecules.

EXAMPLE 6: Preparation of Aryl Derivative of Asp Ala His Lys [SEQ ID NO:1]

A derivative having the formula shown in Figure 1B, wherein R₁ is phenyl, was prepared. Diethylacetamidomalonate (10 gm) in anhydrous ethanol (100 mL) was added to a slurry of sodium ethoxide in ethanol (5 gm; 50 mL) and heated to reflux for 30 min. The product was cooled (10°C) and reacted with ethyl α -bromophenyl acetate (5 gm). The reaction was allowed to proceed to completion (24 h), and excess sodium ethoxide was neutralized with dilute acid. The triester was extracted into ethylacetate and, upon removal of solvent, gave a viscous liquid. The crude product was hydrolyzed with hydrochloric acid (100 mL) and decarboxylated to give phenyl substituted aspartic acid (10 gm). The N-benzoyloxy t-butyl derivative was prepared using a standard reaction sequence. To the resin-bound tripeptide (Lys His Ala) prepared as described in Example 1 (20 mg) in DMF was added the N-benzoyloxy-t-butyl aspartic acid derivative, followed by a mixture of diisopropylamine (8 equivalent) and TBTU- (4 equivalent). The resin was shaken for about 24 h, and the reaction monitored by the ninhydrin test. At the end of this period, DMF was drained, and the resin

was washed with DMF and DCM. The solution was drained, and the beads washed with DCM (3 x 2 ml). The tetrapeptide derivative was isolated by careful hydrolysis. Stereoisomers of the tetrapeptide were separated by preparative-scale HPLC.

EXAMPLE 7: Inhibition Of The Generation Of ROS

By The Tetrapeptide Asp Ala His Lys [SEQ ID NO:1]

A tetrapeptide having the sequence L-Asp L-Ala L-His L-Lys [SEQ ID NO:1] (the L-tetrapeptide) was obtained from one or more companies that provide custom synthesis of peptides, including Ansynth Services, QCB, Genosys and Bowman Research. The peptide was prepared by standard solid phase synthesis methods (see also Example 1).

The ability of the L-tetrapeptide to inhibit the generation of ROS was tested as described in Gutteridge and Wilkins, *Biochim. Biophys. Acta*, **759**, 38-41 (1983) and Cheeseman et al., *Biochem. J.*, **252**, 649-653 (1988). Briefly, Cu(II) and H₂O₂ were mixed causing the generation of hydroxyl radicals in a Fenton-type reaction. The hydroxyl radicals attack the sugar 2-deoxy-D-ribose (the sugar residue of DNA) to produce fragments. Heating the fragments at low pH produces malonaldehyde that, upon the addition of 2-thiobarbituric acid, yields a pink chromogen which can be measured spectrophotometrically at 532 nm. Thus, the absorbance at 532 nm is a measure of the damage to 2-deoxy-D-ribose.

The assay was performed with and without the L-tetrapeptide. The results are summarized in Table 1. As can be seen from Table 1, when the L-tetrapeptide was present at Cu(II):tetrapeptide ratios of 1:1.2 and 1:2, the degradation of 2-deoxy-D-ribose was inhibited by 38% and 73%, respectively. Clearly, the L-tetrapeptide inhibited the degradation of 2-deoxy-D-ribose by hydroxyl radicals.

TABLE 1

	CuCl ₂ (mM)	H ₂ O ₂ (mM)	Tetra-peptide (mM)	OD at 532 nm	Percent Inhibition
Control	0.1	2.0	0.0	0.124	
Tetrapeptide	0.1	2.0	0.12	0.077	38
Control	0.1	2.0	0.0	0.175	
Tetrapeptide	0.1	2.0	0.2	0.048	73

A similar assay was also performed using a tetrapeptide having the sequence Asp Ala His Lys composed of all D-amino acids (D-tetrapeptide). The D-tetrapeptide was obtained from one or more companies that provide custom synthesis of peptides, including Ansynth Services and QCB. The peptide was prepared by standard solid phase synthesis methods (see Example 1)

The ability of the D-tetrapeptide to inhibit the generation of ROS was tested as described by Zhao and Jung, *Free Radic Res*, **23(3)**, 229-43 (1995). Briefly, Cu(II) and ascorbic acid were mixed causing the generation of hydroxyl radicals in a Fenton-type reaction. The advantage of using ascorbic acid instead of hydrogen peroxide is that ascorbic acid does not interfere with other assays (*i.e.* LDH assay) which is not the case with peroxide. The hydroxyl radicals attack the sugar 2-deoxy-D-ribose to produce fragments. Heating the fragments at low pH produces malonaldehyde that, upon the addition of 2-thiobarbituric acid, yields a pink chromogen which can be measured spectrophotometrically at 532 nm. Thus, the absorbance at 532 nm is a measure of the damage to 2-deoxy-D-ribose.

Establishing optimal Cu(II) and ascorbic acid concentrations was the first step in developing this protocol. First, a constant Cu(II) concentration of 10 μ M was used based on this level being the physiological concentration found in the body (bound and unbound Cu(II)). The ascorbic acid concentrations were varied in order to establish a linear range. The ascorbic acid concentration chosen was 500 μ M since it gave the most absorbance at 532 nm and still fell in the linear range. Interestingly, at ascorbic acid concentrations greater than 500 μ M,

there was a steady decrease in hydroxyl radicals presumably due to ascorbic acid's dual effect as a hydroxyl radical generator at low concentrations and an antioxidant at high concentrations.

Using the aforementioned concentrations for Cu(II) and ascorbic acid, a titration curve was established for the D-tetrapeptide. Briefly, the D-tetrapeptide was pre-incubated with Cu(II) for 15 minutes at room temperature prior to adding ascorbic acid. This was done to permit the D-tetrapeptide to bind with the Cu(II) and therefore inhibit ROS generation. As can be seen from the table, when the Cu(II):D-tetrapeptide ratio was between 4:1 to 4:7, there was little to no inhibition of hydroxyl radical generation. When the ratio was 1:2 or higher, there was total inhibition of hydroxyl radical production.

TABLE 2

Cu(II):D-Tetrapeptide	Cu(II) (μM)	Ascorbic Acid (μM)	D-Tetrapeptide (μM)	A532	% Inhibition
1:0	10	500	0	0.767	
4:1	10	500	2.5	0.751	
2:1	10	500	5	0.743	
1:1	10	500	10	0.751	
4:5	10	500	12.5	0.789	
2:3	10	500	15	0.774	
4:7	10	500	17.5	0.737	
1:2	10	500	20	0.029	96.2
1:4	10	500	40	0.016	97.9

**EXAMPLE 8: Testing of Asp Ala His Lys D-Tetrapeptide
In A Langendorff Reperfusion Model**

Blood-perfused hearts were prepared essentially as in previous studies (Galiñanes et al., *Circulation*, **88**:673-683 (1993); Kolocassides et al., *Am. J. Physiol.*, **269**:H1415-H1420 (1995); Hearse et al., *J. Mol. Cell. Cardiol.*, **31**:1961-1973 (1999)). See Figure 8. The procedures are described briefly below.

Male Wistar rats, obtained from Bantin and Kingman Universal, UK, were used. All animals received humane care in compliance with the "*Principles of Laboratory Animal Care*" formulated by the National Society for Medical Research and the "*Guide for the Care and Use of Laboratory Animals*" prepared by the National Academy of Sciences, and published by the US National Institutes of Health (NIH Publication No. 85-23, revised 1996).

Support rats (300-400g) were anesthetized with sodium pentobarbitone (60 mg/kg, intraperitoneally) and anticoagulated with heparin (1000 IU/kg intravenously). The right femoral vein and left femoral artery were exposed by blunt dissection and cannulated (18G and 22G Abbocath-T catheters respectively) for the return and supply of blood to the perfused heart. An extracorporeal circuit was established, primed with Gelofusine® plasma substitute (B. Braun Medical Ltd., Aylesbury, UK) and was maintained for 15 minutes (min) before connection to the isolated heart. This period was to ensure that the priming solution was adequately mixed with the blood of the support rat and that the entire preparation was stable. Each 500 ml of Gelofusine® contains 20.00 g succinated gelatin (average molecular weight 30,000), 3.65 g. sodium chloride, water for injection to 500 ml (electrolytes mmol/500 ml: cations Na 77, anions Cl 62.5, pH 7.4). Prior to perfusing the donor heart, an additional 7-8 ml of blood from a rat of the same strain was added to the central reservoir. This was to ensure that the support rat had an adequate supply of blood during the experiment when blood was not recirculated but instead collected for a 2 min period. A peristaltic pump (Gilson Minipuls 3) was located on the arterial outflow of the support rat and flow through the extracorporeal circuit was increased gradually over 10 min to a value of 2.5 ml/min. This gradual increase prevented the drop in arterial pressure that would have occurred if a flow rate of 2.5 ml/min had been established immediately. The blood was pumped through a cannula (to which the aorta of the perfused heart would subsequently be attached) and returned, by gravity, via a reservoir and filter to the venous inflow line of the support animal. An air-filled syringe above the perfusion cannula acted as a compliance chamber, which served to dampen oscillations in perfusion pressure which occurred as a consequence of the contraction of the isolated heart and the peristaltic action of the pump. The support animal was allowed to

breathe a mixture of 95% O₂ + 5% CO₂ through a 35% Venturi face mask. The flow rate was adjusted to maintain blood pO₂ and pCO₂ within the physiological range. Body temperature was stabilized at 37.0 (±0.5) °C by means of a thermostatically-controlled heating pad and was monitored by a rectal thermometer. Blood pressure was monitored by means of a pressure transducer attached to the arterial line. All pressure transducers were connected to a MacLab (ADInstruments, Australia), which was run continuously through the experiment. Blood gas (pO₂, pCO₂, pH), hematocrit, glucose and electrolyte levels (Na⁺, K⁺, Ca²⁺) of the support rat were monitored before the donor heart was attached to the extracorporeal circuit and at the end of each experiment. During the course of the experiment, minimum amounts of donor blood (from another rat of the same strain) were transfused as required so as to maintain the volume and stability of the preparation. Additional heparin and pentobarbitone were administered into the central reservoir as required.

To isolate hearts, each rat (270-350 g) was anesthetized with diethyl ether and anticoagulated with heparin (1000 IU/kg intravenously). The heart was then immediately excised and immersed in cold (4°C) Gelofusine®. The aorta was rapidly cannulated and perfused in the Langendorff mode, (Langendorff, *Pflügers Archives für die Gesamte Physiologie des Menschen und der Tiere*, 61:291-332 (1895)) with arterial blood from the support animal, at a constant flow rate of 2.5 ml/min. After removal of the left atrial appendage, a fluid-filled balloon catheter (for the measurements of left ventricular systolic and diastolic pressures, and, by difference, left ventricular developed pressure), attached to a pressure transducer, was introduced into the left ventricle via the mitral valve. The balloon was inflated with water until a left ventricular end diastolic pressure (LVEDP) of between 4-8 mmHg was obtained. Heart rate was calculated from the pressure trace and expressed as beats per minute (bpm). Perfusion pressure was measured via a sidearm of the aortic cannula. All pressure transducers were connected to a MacLab, which was run continuously through the experiment.

Excised hearts were randomly assigned to two treatment groups (see Figure 9; 6 hearts/group) and aerobically perfused for 20 min prior to (i) saline control with 2 min saline

infusion immediately prior to a 30 min period of ischemia plus 2 min saline infusion at the onset of reperfusion and (ii) drug with 2 min drug infusion immediately prior to a 30 min period of ischemia (the drug was therefore trapped in the vasculature for the duration of the ensuing ischemic period), plus a 2 min drug infusion at the onset of reperfusion. Hearts were then subjected to 30 min of global, zero-flow ischemia, during which time they were immersed in saline at 37.0°C. Ischemia was initiated by clamping the line leading from the pump to the aortic cannula, thus diverting the flow away from the isolated heart back to the support animal, via the bypass line. Hearts were then reperfused for 40 min, during which time contractile function was continuously measured.

The drug, whose identity was unknown to the researchers performing the experiments, was the tetrapeptide D-Asp D-Ala D-His D-Lys. The tetrapeptide was supplied to the researchers by Bowman Research, UK, dissolved in saline at a concentration of 16.7 mg/mL. It was infused as supplied without any dilution or modification. Physiological saline was supplied by Baxter, UK and used in controls. Fresh solutions of saline and the drug were used daily.

Drug or vehicle was infused into a sidearm of the aortic cannula by means of a peristaltic pump (Gilson Minipuls 3), set at a constant flow of 0.25 ml/min. Since blood flow through the aortic cannula was 2.5 ml/min, and drug infusion was 0.25 ml/min, the final concentration of drug delivered to the heart was 1/11th of that supplied by Bowman Research. During the 2 min period of pre-ischemic vehicle or drug infusion and at the time points indicated in Figure 9 arterial and venous blood samples were collected, centrifuged and frozen for analysis. The infusion was then repeated for the first 2 min of the reperfusion period, during which time the blood was not collected, but recirculated.

Predefined exclusion criteria stated that: (i) support animals would be excluded from the study if they did not attain a stable systolic blood pressure ≥ 80 mm Hg before cannulation of the donor heart, (ii) donor hearts would be excluded from the study if, at the 20 min baseline pre-intervention reading, LVDP ≤ 100 mm Hg or (iii) blood chemistry values were outside the normal range.

Results are expressed as mean \pm SEM. All recovery values are expressed as a percent of the pre-intervention baseline value (measured 20 min after the onset of the experiment) for each individual heart. The two-tailed unpaired Student's *t* test was used for the comparison of two means between groups. A difference was considered statistically significant when $p < 0.05$.

For reasons of quality control and to allow application of predefined exclusion criteria, the stability and reproducibility of the system were monitored by measuring the blood chemistry (pH, pO₂, pCO₂, haematocrit, Na⁺, K⁺ and Ca²⁺, glucose) and baseline contractile function of each support animal (immediately before perfusing a donor heart and at the end of each experiment) and each perfused heart. Table 3 reveals that there were only minor changes in each index measured, confirming that similar perfusion conditions applied in both study groups and that all values were within the acceptable physiological range. The systolic pressure and heart rate of the support rats are shown in Table 4. As can be seen, there were no significant differences between the two study groups at the 15 min baseline reading.

Table 5 shows that there were no significant differences between groups at the end of the 20 min aerobic perfusion period (*i.e.* just prior to the infusion of drug or vehicle) in LVDP, heart rate and perfusion pressure. Thus, for LVDP, the primary endpoint in the study, the mean values were 177.3 ± 10.6 mmHg and 177.2 ± 5.6 mmHg for the groups that were to become saline control and drug treated.

As expected, myocardial ischemia caused cessation of myocardial contraction, with cardiac arrest initially in the diastolic state. However, as ischemic injury developed with time an increase in diastolic state occurred as the heart went into ischemic contracture. The temporal profiles for the development of ischemic contracture in each of the study groups is shown in Figure 10. As can be seen from Table 6, there were no differences in any of the measured indices, although there is some evidence of a trend to delay time-to-50% contracture in the drug treated group, which is suggestive with protection.

Figure 11 shows the profiles for the mean recovery of LVDP (expressed as a percent of baseline pre-intervention values) in both study groups. It is evident that hearts in the saline

control group recovered slowly and poorly, such that by the end of the 40 min reperfusion period, LVDP was only $15.3 \pm 3.2\%$ of the pre-intervention control. By contrast, hearts in the drug group recovered more rapidly and to a greater extent ($50.5 \pm 9.3\%$).

Figure 12 shows the absolute values for the left ventricular end diastolic pressure in both study groups during the 40 min period of reperfusion. In both groups, the high levels of LVEDP resulting from the contracture which developed during ischaemia fell with time towards the pre-intervention control value. However, the drug group normalized their LVEDP more quickly and more completely than that seen in the saline control group, the difference being significant at every time point studied. This would be consistent with the enhanced recovery seen during reperfusion in the drug group.

A comparison of the heart rates obtained in the saline control group and drug, as shown in Figure 13 reveals that these two groups were essentially identical (115.0 ± 3.8 versus $127.2 \pm 22.8\%$).

As shown in Figure 14, the perfusion pressure during the 40 min reperfusion period was essentially constant in both groups and the values did not differ significantly between each other at the end of the 40 min reperfusion period. In saline controls, the mean value at the end of reperfusion was $109.9 \pm 8.2\%$ of its pre-intervention control, while that of the drug group was $87.4 \pm 8.0\%$ of its pre-intervention value. Thus, although the values for drug tended to be lower throughout the reperfusion period (which is consistent with the observed cardioprotection), these changes did not reach statistical significance.

The results of this pilot study indicate that, in the isolated blood-perfused rat heart, Asp Ala His Lys appears to have significant and substantial protective properties as assessed by an approximately three and a half (3.5) fold ($15.3 \pm 3.2\%$ to $50.5 \pm 9.3\%$) enhancement of post-ischemic functional recovery. The magnitude of protection is equal to some of the most powerful interventions studied.

TABLE 3
Composition of the blood perfusing the isolated donor hearts

Index	Value prior to attaching donor heart		End of experiment	
	Saline Control	Drug*	Saline Control	Drug*
pH	7.29±0.01	7.26±0.01	7.33±0.02	7.31±0.02
pCO ₂ (mmHg)	60.0±3.2	67.7±2.2	57.6±4.9	65.6±4.7
pO ₂ (mmHg)	230.6±14.8	281.7±23.2	241.0±37.2	252.3±33.9
Haematocrit (%)	27.5±0.9	27.3±1.8	26.8±1.7	27.0±1.5
Na ⁺ (mmol/L)	146.7±0.3	146.2±0.6	145.7±0.2	146.2±0.5
K ⁺ (mmol/L)	3.3±0.1	3.2±0.1	4.2±0.2	4.2±0.1
Ca ²⁺ (mmol/L)	1.2±0.1	1.2±0.1	1.4±0.1	1.4±0.1
Glucose (mmol/L)	8.9±0.6	10.2±0.3	10.2±0.5	10.2±0.7
% oxygen saturation	99.6±0.1	99.8±0.1	98.7±1.2	99.7±0.1

There were 4-6 support animals per group. All values are expressed as mean ± SEM.

* Drug was D-Asp D-Ala D-His D-Lys.

TABLE 4

Baseline systolic pressure and heart rate in the support blood-perfused rat

Subsequent Treatment Group	Control values (t=15 min aerobic perfusion)	
	Systolic Pressure (mmHg)	Heart Rate (bpm)
Saline Control	96.4±2.6	296.6±9.8
Drug*	98.1±3.8	297.6±8.7

All values are expressed as mean ± SEM. There were 6 support animals per group.

* Drug was D-Asp D-Ala D-His D-Lys.

TABLE 5

Baseline left ventricular developed pressure (LVDP), heart rate and perfusion pressure in isolated blood-perfused rat hearts *before* various interventions

Subsequent Treatment Group	Control values (t=20 min aerobic perfusion)		
	LVDP (mmHg)	Heart Rate (bpm)	Perfusion Pressure (mmHg)
Saline Control	177.3±10.6	236.6±17.9	94.6±7.5
Drug*	177.2±5.6	257.3±29.7	99.6±7.3

All values are expressed as mean ± SEM. There were 6 support animals per group.

* Drug was D-Asp D-Ala D-His D-Lys.

TABLE 6

Ischemic contracture during 30 min of global, zero flow ischaemia

Group	Contracture			
	Initiation (min)	Time-to-50% (min)	Peak (mmHg)	Time-to-peak (min)
Saline Control	9.2±1.6	15.1±0.5	93.7±3.0	19.8±0.6
Drug*	11.9±1.7	16.5±0.9	89.5±3.0	21.7±1.2

All values are expressed as the mean ± SEM. There were 6 hearts per group.

* Drug was D-Asp D-Ala D-His D-Lys.

EXAMPLE 9: Testing of Asp Ala His Lys In A Brain Ischemia Model

Focal ischemic infarcts were made in mature male Wistar rats (270-300 g, Charles River Laboratories) as described previously (Koizumi et al, *Jpn. J. Stroke*, **8**:1-8 (1986); Chen et al., *J. Cereb.lood Flow Metab.*, **12**(4):621-628 (1992)). Animals were allowed free access to food and water before surgery. They were anesthetized with 3.5% halothane, and anesthesia was maintained with 1.0% - 2.0% halothane in 70% N₂/30% O₂ using a face mask. Rectal temperature was maintained at 37°C during surgery using a feedback-regulated water heating system (YSI 73A rectal probe, Fisher, connected to a K-20/64N aquatic blanket, Hamilton Industries, Cincinnati, OH). Previous studies have shown that rectal and brain

temperatures are identical during and after ischemia in this model (Chen et al., *J. Cereb.lood Flow Metab.*, **12**(4):621-628 (1992)). The right femoral artery was cannulated with medical grade silicone tubing (Technical Products, Inc., Decatur, GA) for measurement of blood gases and blood pressure, and the femoral vein was cannulated for infusions. Cannulae were drawn through a subcutaneous tunnel and exited through the dorsal neck. Arterial blood gases were measured in all animals before and after ischemia.

For ischemia surgery, the right common carotid artery was exposed at its bifurcation. A 4-0 nylon suture, with its tip rounded by heating over a flame, was then advanced 18.5-19.5 mm (depending on the animal's weight) from the external into the internal carotid artery and then through the intracranial carotid artery until the tip occluded the origin of the middle cerebral artery (MCA). Animals were then allowed to awaken from anesthesia. At 2 hours after MCA occlusion, they were re-anaesthetized, and intra-arterial sutures were withdrawn into the external carotid artery.

Beginning one minute prior to occlusion, animals received an intravenous infusion of vehicle alone (control) or drug (D-Asp D-Ala D-His D-Lys) in vehicle over one minute. The identity of the drug was unknown to the researchers performing the experiments. It was supplied to the researchers as a concentrated stock (16.67 mg/ml) in phosphate buffered saline, pH 7.4, and was stored it at -80°C. The drug was determined to be biologically active prior to use by determining its ability to reduce free radical formation *in vitro* as described in Example 7. The stock solution was thawed just prior to use, and a sufficient quantity was administered to give a dose of 20 mg/kg. At the end of the intravenous administration of the drug or vehicle, the nylon suture was immediately advanced to occlude the MCA. Following the 2 hours of occlusion of the MCA, the animals received a repeat intravenous infusion of drug or vehicle over one minute. At the end of the second infusion, the nylon suture was immediately pulled back from occluding the MCA to allow for reperfusion. Also, after the second infusion, the animals were re-anesthetized, and the cannulae were removed. The animals were returned to their home cages, where they were allowed free access to food and water.

Animals were weighed before ischemia and before sacrifice. A neurological examination, as described in Zea Longa et al., *Stroke*, **20**:84-91 (1989), was administered at 1 hour and at 24 hours after reperfusion. Scoring was as follows: 0, normal; 1, failure to extend contralateral (left) forepaw fully (a mild focal neurologic deficit); 2, circling to the left (a moderate focal neurologic deficit); 3, falling to the left (a severe focal neurologic deficit); and 4, no spontaneous gait and depressed level of consciousness.

Twenty-four hours after MCA occlusion, animals were anesthetized with ketamine (44 mg/kg) and xylazine (13 mg/kg), both given intramuscularly, and perfused transcardially with heparinized saline, followed by 10% buffered formalin. The brains were removed and cut into 2-mm coronal slices using a rat brain matrix (Activational System, Inc., Warren, MI; a total of 7 slices). The slices were then embedded in paraffin, and 6-mm sections were cut from the anterior surface of each slice and stained with hematoxylin and eosin (H and E). Infarct volume was determined using a computer-interfaced image analysis system (Global Lab Image system, Data Translation, Marlboro, MA), using the "indirect" method (Swanson et al., *J. Cerebral Blood Flow Metabol.*, **10**:290-293 (1990)): the area of intact regions of the ipsilateral (right) hemisphere and area of the intact contralateral (left) hemisphere were determined for each slice, the former was subtracted from the latter to calculate infarct area per slice. Infarct areas were then summed and multiplied by slice thickness to yield infarct volume per brain (in mm³).

The results are presented in Tables 7-10 below. Some of the data are expressed as mean \pm S.E.M. Continuous data were analyzed by repeated measures ANOVA and paired or unpaired two-tailed t-tests with Bonferroni correction where appropriate. Non-continuous behavior data were analyzed by the Mann-Whitney U-test.

TABLE 7
Infarct Volume

Treated*		Control	
Animal	Infarct Volume (mm ³)	Animal	Infarct Volume (mm ³)
#1	11.5	#2	44.5
#3	14.7	#4	32.3
#5	43.3	#6	39.4
#7	10.9	#8	22.2
Mean	20.1	Mean	34.6
S.E.M.	7.7	S.E.M.	4.8

* Treated with D-Asp D-Ala D-His D-Lys

TABLE 8
Neurological Scale

Treated*			Control		
Animal	Day 0 ^a	Day 1 ^b	Animal	Day 0 ^a	Day 1 ^b
#1	2	1	#2	2	2
#3	2	1	#4	2	2
#5	2	2	#6	2	2
#7	2	1	#8	2	2

* Treated with D-Asp D-Ala D-His D-Lys

^a Day of ischemia surgery

^b One day after ischemia surgery

TABLE 9
Body Weight

Treated*			Control		
Animal	Day 0 ^a	Day 1 ^b	Animal	Day 0 ^a	Day 1 ^b
#1	300	272	#2	300	251
#3	300	266	#4	300	256
#5	295	256	#6	278	230
#7	295	255	#8	300	250

* Treated with D-Asp D-Ala D-His D-Lys

^a Day of ischemia surgery

^b One day after ischemia surgery

TABLE 10

Blood Gases And Blood Pressure

10 minutes before MCA occlusion					10 minutes after MCA occlusion				
Animal	pH	pCO ₂	pO ₂	BP ^c	Animal	pH	pCO ₂	pO ₂	BP ^c
#1	7.437	38.5	117.1	96	#1	7.422	40.2	130.1	106
#3	7.430	38.3	110.5	89	#3	7.423	40.1	130.1	98
#5	7.518	38.5	147.0	98	#5	7.471	36.5	105.3	92
#7	7.423	33.4	97.4	86	#7	7.433	34.0	139.9	89
#2	7.401	35.5	120.1	90	#2	7.423	40.1	130.1	103
#4	7.440	38.5	121.1	93	#4	7.421	39.5	129.3	101
#6	7.425	35.9	110.0	88	#6	7.453	36.2	105.3	92
#8	7.417	39.5	120.7	90	#8	7.428	36.9	111.0	93

EXAMPLE 10: Inhibition Of The Generation Of ROS



The ability of the tetrapeptide L-Asp L-Ala L-His L-Lys [SEQ ID NO:1] and other peptides and compounds to inhibit the production of ROS was tested. The other peptides tested were: L-Asp L-Ala L-His L-Lys L-Ser L-Glu L-Val L-Ala L-His L-Arg L-Phe L-Lys [SEQ ID NO:3]; L-Ala L-His L-Lys L-Ser L-Glu L-Val L-Ala L-His L-Arg L-Phe L-Lys [SEQ ID NO:4]; L-His L-Lys L-Ser L-Glu L-Val L-Ala L-His L-Arg L-Phe L-Lys [SEQ ID NO:5]; and Acetylated-L-Asp L-Ala L-His L-Lys L-Ser L-Glu L-Val L-Ala L-His L-Arg L-Phe L-Lys [SEQ ID NO:6]. The peptides were obtained from one or more companies that provide custom synthesis of peptides, including Ansynth Services, QCB, Genosys and Bowman Research. The other compounds tested were histidine (Sigma Chemical Co.), catalase (Sigma Chemical Co.), and superoxide dismutase (Sigma Chemical Co.).

1. Inhibition Of Hydroxyl Radical Production

The hydroxyl radical is probably the most reactive oxygen-derived species. The hydroxyl free radical is very energetic, short-lived and toxic.

Some researchers suggest that the toxicity of hydrogen peroxide and superoxide radical may be due to their conversion to the hydroxyl free radical. The superoxide radical can be directly converted to the hydroxyl radical via the Haber-Weiss reaction. Alternatively, it can be converted to hydrogen peroxide which, in turn, is converted into the hydroxyl radical via the Fenton reaction. Both pathways require a transition metal, such as copper (Acworth and Bailey, *The Handbook Of Oxidative Metabolism* (ESA, Inc. 1997)).

It is also known that copper, in the presence of ascorbate, produces hydroxyl radicals. The following reaction scheme has been suggested:



Biaglow et al., *Free Radic. Biol. Med.*, **22**(7):1129-1138 (1997).

The ability of the compounds listed above to inhibit the generation of hydroxyl radicals was tested as described in Gutteridge and Wilkins, *Biochim. Biophys. Acta*, **759**:38-41 (1983). Briefly, Cu(II) and ascorbic acid were mixed causing the generation of hydroxyl radicals. Then, deoxyribose was added, and the hydroxyl radicals, if present, attacked the deoxyribose to produce fragments. Heating the fragments at low pH produced malonaldehyde that, upon the addition of 2-thiobarbituric acid (TBA), yielded a pink chromogen which was measured spectrophotometrically at 532 nm. Thus, absorbance at 532 nm is a measure of the damage to deoxyribose and, therefore, of hydroxyl radical formation.

To perform the assay, CuCl₂ in buffer (20 mM KH₂PO₄ buffer, pH 7.4) and either one of the test compounds in buffer or buffer alone were added to test tubes (final concentration of CuCl₂ was 10 μM). The test tubes were incubated for 15 minutes at room temperature. Then, 0.5 mM ascorbic acid in buffer and 1.9 mM 2-deoxy-D-ribose in buffer were added to each test tube, and the test tubes were incubated for 1 hour at 37°C. Finally, 1 ml of 1% (w/v) TBA in 50 mM NaOH and 1 ml of concentrated acetic acid were added to each test tube, and the test tubes were incubated in boiling water for 15 minutes. After the test tubes had cooled for 15 minutes, the absorbance at 532 nm was read.

It was found that the tetrapeptide L-Asp L-Ala L-His L-Lys [SEQ ID NO:1] caused complete inhibition of the formation of hydroxyl radicals in this assay at tetrapeptide/copper ratios of 2:1 or higher. Tetrapeptide/copper ratios less than 2:1 were ineffective.

The results of a time course are presented in Figure 15A. As can be seen in Figure 15A, copper and ascorbate (no added peptide) produced TBA-reactive substances quickly and reached a maximum in 30 minutes. The tetrapeptide at a tetrapeptide/copper ratio of 2:1 prevented all formation of TBA-reactive substances. Interestingly, the tetrapeptide at a tetrapeptide/copper ratio of 1:1 slowed the production of TBA-reactive substances. These data suggest that the tetrapeptide at a 1:1 tetrapeptide/copper ratio is able to offer some protection from hydroxyl radicals by binding copper which results in site-directed hydroxyl attack on the tetrapeptide. Once enough of the tetrapeptide is destroyed, then copper is released, which allows it to produce hydroxyl radicals that attack the deoxyribose.

When the tetrapeptide at a tetrapeptide/copper ratio of 2:1 was incubated for longer periods of time, its ability to prevent the formation of TBA-reactive substances slowly eroded. See Figure 15B. As can be seen from Figure 15B, the production of TBA-reactive substances was inhibited by 95% during the first 4 hours of incubation. By 24 hours, the level of inhibition had dropped to 50% and, by 48 hours, the level of inhibition had dropped to 20%. These data suggest that TBA-reactive substances are still being produced even in the presence of the tetrapeptide. They also suggest that the tetrapeptide is being degraded during the time course of the experiment. This degradation is more than likely due to the formation of free radicals in close proximity to the tetrapeptide/copper complex which attack and degrade the tetrapeptide, with release of the copper. Since free radicals, such as the hydroxyl radical, are very reactive, they will attack the first electron rich molecule they come into contact with, which would be the tetrapeptide in this case.

The effect of pH on the inhibition of hydroxyl radical formation by the tetrapeptide was tested at a tetrapeptide/copper ratio of 2:1. At this ratio, the tetrapeptide gave >95% inhibition of the formation of TBA-reactive species at pH 7.0-8.5. These are physiological pH levels and pH levels that would be expected during ischemia (acidosis occurs in ischemic tissues). At pH 6.0, the tetrapeptide was ineffective at preventing the formation of TBA-reactive species, possibly due to the reduced ability of the histidine to bind copper. The nitrogen atom on the imidazole ring of histidine participates in binding copper with a pKa of 6.0. Therefore, at a pH of 6.0, histidine is only able to bind 50% of the copper. The other 50% of the copper would be unbound or loosely bound to the tetrapeptide by the other amino acids and would, therefore, be able to participate in the production of TBA-reactive species.

Histidine and several peptides with histidine in different positions were tested at 1:1 and 2:1 peptide:copper ratios for their ability to inhibit the production of hydroxyl radicals. Also, a peptide having an acetylated aspartic acid (Ac-Asp) as the N-terminal amino acid was also tested. The results are presented in Table 11. In Table 11, the % inhibition is the percent decrease in absorbance compared to buffer alone divided by the absorbance of the buffer alone.

As can be seen from the results in Table 11, the peptides with histidine in the second and third positions gave >95% inhibition at a 2:1 peptide:copper ratio, while these peptides at a 1:1 peptide:copper ratio were ineffective. Interestingly, at a 2:1 peptide:copper ratio, the peptide with histidine in the first position and the peptide with acetylated aspartic acid as the N-terminal amino acid provided some protection (about 47% and about 28% inhibition, respectively), although this protection might be attributable to the histidine in the seventh and ninth positions, respectively, of these peptides. Histidine alone at a 2:1 histidine:copper ratio provided some protection (about 20% inhibition).

Catalase has been shown to prevent hydroxyl radical formation. Gutteridge and Wilkins, *Biochim. Biophys. Acta*, **759**:38-41 (1983); Facchinetti et al., *Cell. Molec. Neurobiol.*, **18**(6):667-682 (1998); Samuni et al., *Eur. J. Biochem.*, **137**:119-124 (1983). Catalase (0-80 nM) was, therefore, tested in this assay, and it was found to prevent the formation of the pink chromogen (data not shown). This finding suggests that hydrogen peroxide is formed in this assay, since catalase breaks down hydrogen peroxide to water and agrees with Equations 3 and 4 above. Catalase also prevents the formation of the pink chromogen when the L-Asp L-Ala L-His L-Lys [SEQ ID NO:1] tetrapeptide at a tetrapeptide/copper ratio of 1:1 is present (data not shown). As shown above, at this ratio, the copper is still able to participate in the redox reactions to produce hydroxyl radicals. These experiments show that hydrogen peroxide is an important precursor to the formation of the hydroxyl radical.

TABLE 11

Compound (Ratio) ^a	Absorbance at 532 nm	Absorbance at 532 nm	% Inhibition
Copper only (buffer control)	0.767*	0.954	0
Histidine/copper (2:1)		0.760	20.3
His Lys Ser Glu Val Ala His Arg Phe Lys ^b /copper (1:1)		0.716	24.9
His Lys Ser Glu Val Ala His Arg Phe Lys ^b /copper (2:1)		0.509	46.6
Ala His Lys Ser Glu Val Ala His Arg Phe Lys ^c /copper (1:1)		0.843	11.6
Ala His Lys Ser Glu Val Ala His Arg Phe Lys ^c /copper (2:1)		0.047	95.1
Asp Ala His Lys Ser Glu Val Ala His Arg Phe Lys ^d /copper (1:1)	0.645		13.2
Asp Ala His Lys Ser Glu Val Ala His Arg Phe Lys ^d /copper (2:1)	0.040		95.8
Ac-Asp Ala His Lys Ser Glu Val Ala His Arg Phe Lys ^e /copper (1:1)	0.633		16.9
Ac-Asp Ala His Lys Ser Glu Val Ala His Arg Phe Lys ^e /copper (2:1)	0.692		27.5
Asp Ala His Lys ^f /copper (1:1)	0.751*		1.3
Asp Ala His Lys ^f /copper (2:1)	0.029*		96.2

^a All amino acids are L-amino acids.

^b SEQ ID NO:5

^b SEQ ID NO:5

^c SEQ ID NO:4

^c SEQ ID NO:4

^d SEQ ID NO:3

^d SEQ ID NO:3

^e SEQ ID NO:6

^e SEQ ID NO:6

^f SEQ ID NO:1

^f SEQ ID NO:1

* Data taken from Table 2 in Example 7.

B. Assay For Superoxide Dismutase (SOD) Activity

The enzyme superoxide dismutase (SOD) is a naturally-occurring enzyme which is responsible for the breakdown in the body of superoxide to hydrogen peroxide (similar to Equation 3). Hydrogen peroxide can then be detoxified by catalase.

SOD was assayed for activity in the assay described in the previous section and was found to have none (data not shown). This result is not surprising since SOD actually converts superoxide radical into hydrogen peroxide. Hydrogen peroxide can then be converted into the hydroxyl radical by reduced copper.

There are reports in the literature that copper complexes have SOD activity. Athar et al., *Biochem. Mol. Biol. Int.*, **39**(4):813-821 (1996); Ciuffi et al., *Pharmacol Res.*, **38**(4):279-287 (1998); Pogni et al., *J. Inorg. Biochem.*, **73**:157-165 (1999); Willingham and Sorenson, *Biochem. Biophys. Res. Commun.*, **150**(1):252-258 (1988); Konstantinova et al., *Free Rad. Res. Comms.*, **12-13**:215-220 (1991); Goldstein et al., *J. Am. Chem. Soc.*, **112**:6489-6492 (1990). This finding is not surprising since SOD itself has copper in its active site.

The SOD activity of copper complexes of the tetrapeptide L-Asp L-Ala L-His L-Lys [SEQ ID NO:1] was assayed. Superoxide radicals were produced using the xanthine oxidase assay of Beauchamp and Fridovich, *Anal. Biochem.*, **44**:276-287 (1971). Xanthine oxidase converts xanthine into uric acid, with oxygen acting as an electron acceptor. This causes superoxide radical to be produced. Superoxide radical is able to reduce nitro blue tetrazolium (NBT). Reduced NBT has a λ_{max} of 560 nm. It is known that copper inhibits xanthine oxidase activity (Konstantinova et al., *Free Rad. Res. Comms.*, **12-13**:215-220 (1991)), so all experiments containing copper also contained ethylenediaminetetracetic acid (EDTA), a known copper chelator. The EDTA-copper complex was tested for SOD activity and was shown to have no SOD activity (data not shown).

To perform the assay for SOD activity, 0.1 mM xanthine (Sigma Chemical Co.), 25 μ M NBT (Sigma Chemical Co.), 50 mM sodium carbonate, and 1.2 μ M EDTA (Sigma Chemical Co.), were mixed in a cuvette (all final concentrations, final pH 10.2). The reaction was started by the addition of various amounts of a tetrapeptide-copper complex

(tetrapeptide/copper ratios of 1:1 and 2:1) and 20 nM xanthine oxidase (Sigma Chemical Co.). The tetrapeptide-copper complex was prepared by mixing the tetrapeptide and copper (as CuCl_2) and allowing the mixture to incubate for 15 minutes at room temperature immediately before addition to the cuvette. The samples were read at time 0 and every 60 seconds for five minutes at 560 nm.

The complex of the tetrapeptide with copper at a ratio of 1:1 was shown to have SOD activity, as evidenced by inhibition of NBT reduction (see Figure 16). However, the complex was about 500 times less effective than SOD itself, based on IC_{50} values (amount that gives 50% inhibition) in this assay. The complex of the tetrapeptide with copper at a ratio of 2:1 was found to have no SOD activity (data not shown).

To verify that the 1:1 tetrapeptide-copper complex did not interfere with xanthine oxidase activity, uric acid production was measured at 295 nm. Athar et al., *Biochem. Mol. Biol. Int.*, **39**(4):813-821 (1996); Ciuffi et al., *Pharmacol Res.*, **38**(4):279-287 (1998). This assay is similar to the SOD assay, except that NBT is not present. Instead, uric acid is assayed at 295 nm every 60 seconds for 5 minutes. It was found that the 1:1 tetrapeptide-copper complex only inhibited uric acid production by 11% at a concentration of 600 nM (data not shown). Therefore, the 1:1 tetrapeptide-copper complex has true SOD activity. Since superoxide is converted to hydrogen peroxide by the complex, this could help to explain why it is not effective at preventing hydroxyl radical production.

Superoxide radical production was measured in solutions containing the 1:1 or 2:1 tetrapeptide-copper complexes. The assay combined techniques from the TBA assay and the xanthine oxidase assay. NBT was added to all test tubes in order to quantitate its reduction by superoxide radical. The samples also contained ascorbate and copper and were incubated at 37°C. At 5, 15, 30 and 60 minutes, the samples were removed from the incubator and read at 560 nm. The results are shown in Figure 17. In the sample containing the 2:1 tetrapeptide-copper complex, NBT reduction increased over time and reached a maximum at 30 minutes. The sample containing the 1:1 tetrapeptide-copper complex also showed an increase in NBT reduction, with a decreased maximum reached at 60 minutes. These data suggest that

superoxide accumulates in the sample containing the 2:1 tetrapeptide-copper complex, while the 1:1 tetrapeptide-copper complex mimics superoxide dismutase.

The likely sequence of events that occurs in the production of hydroxyl radicals is as follows:



It has already been shown that the 1:1 tetrapeptide-copper complex can convert superoxide radical ($\text{O}_2^{\cdot -}$) into hydrogen peroxide (H_2O_2). This is the SOD activity of the complex. The 2:1 tetrapeptide-copper complex cannot facilitate this conversion since the two molecules of the tetrapeptide fill all six coordination bonds of copper. This explains why the 2:1 tetrapeptide-copper complex is so effective because it inhibits the formation of hydrogen peroxide, which could in turn react with reduced copper to produce hydroxyl radicals via the Fenton reaction. The 1:1 tetrapeptide-copper complex also provides a valuable service by eliminating the superoxide radical. Even though it produces hydrogen peroxide, most compartments of the human body have sufficient quantities of the enzyme catalase that can eliminate hydrogen peroxide. In the brain, however, catalase activity is reported to be minimal. Halliwell et al., *Methods in Enzymol.*, **186**:1-85 (1990). Therefore, the brain is a particularly vulnerable organ during periods of ischemia, since copper is released due to the acidosis that accompanies ischemia.

C. Protection of DNA

DNA strand breaks were measured according to the method of Asaumi et al., *Biochem. Mol. Biol. Int.*, **39**(1):77-86 (1996). Briefly, 17 $\mu\text{g/ml}$ of plasmid pBR322 DNA was allowed to pre-incubate for 15 minutes at room temperature with 50 μM CuCl_2 and concentrations of the tetrapeptide of 0-200 μM . Then, 2.5 mM ascorbate was added to each reaction, and the mixture was incubated for 1 hour at 37°C. The total volume of the mixture was 16 μL . Next, 3 μL of loading buffer containing 0.25% (w/v) bromophenol blue, 0.25% (w/v) xylene cyanole FF, and 40% (w/v) sucrose in water was added. The samples were separated by electrophoresis in a 0.8% agarose gel for 90 minutes at 70 Volts. The gel was stained in 1X

[illegible]

65